

# Relationship between childhood maltreatment and geriatric depression: the mediator effect of personality traits

Gabriel Behr Gomes Jardim,<sup>1</sup> Armin von Gunten,<sup>2</sup> Irenio Gomes da Silva Filho,<sup>1</sup> Patricia Klarmann Ziegelmann,<sup>3</sup> Daniela Benzano Bumaguin,<sup>3</sup> Eduardo Lopes Nogueira,<sup>1</sup> Paula Engroff,<sup>1</sup> and Alfredo Cataldo Neto<sup>1,4</sup>

<sup>1</sup>Instituto de Geriatria e Gerontologia - IGG, Pontifícia Universidade Católica do Rio Grande do Sul - PUCRS, Brazil

<sup>2</sup>Service Universitaire de Psychiatrie de l'Age Avance - SUPAA, University of Lausanne - UNIL, Switzerland

<sup>3</sup>PPG Epidemiologia e Departamento de Estatística da Universidade Federal do Rio Grande do Sul - UFRGS, Brazil

<sup>4</sup>Núcleo Específico de Formação em Psiquiatria, Escola de Medicina da Pontifícia Universidade Católica do Rio Grande do Sul, Brazil

## ABSTRACT

**Background:** Childhood maltreatment is an important factor associated with adverse mental health outcomes including geriatric depression and the “big five” personality characteristics. The objective of this study was to evaluate a model where personality characteristics mediate the relationship between childhood maltreatment and geriatric depression.

**Method:** In this cross-sectional study, elderly subjects from socioeconomically disadvantaged neighborhoods of Porto Alegre, Brazil ( $n = 260$ ) completed the Childhood Trauma Questionnaire (CTQ), NEO-Five Factor Inventory (NEO-FFI), and Mini International Neuropsychiatric Interview 5.0 (MINI plus). We used structural equation modeling (SEM) to evaluate the mediation hypothesis.

**Results:** The five personality factors (neuroticism, extraversion, agreeableness, openness, and conscientiousness) were related to childhood maltreatment and depression. Mediation analysis revealed that neuroticism and extraversion are complete mediators, agreeableness and conscientiousness are partial mediators, and openness is not a mediator.

**Conclusions:** These findings support the hypothesis in which childhood maltreatment is associated with geriatric depression and mediated by personality factors. These results suggest that reducing the maladaptive personality trait in elderly people who suffered childhood maltreatment could prevent geriatric depression.

**Key words:** childhood maltreatment, geriatric depression, personality traits, elderly

## Introduction

Depression is one of the most frequent mental diseases in the elderly (Kuhlman *et al.*, 2013; Peninx *et al.*, 1999; Raposo *et al.*, 2014). It is associated with worse outcomes from clinical diseases, worse quality of life (Dube *et al.*, 2003), and increased suicide risk (Conwell *et al.*, 2002; Sachs-Ericsson *et al.*, 2016). The city of Porto Alegre, Brazil has a high prevalence (30.6%) of geriatric depression among the socioeconomically disadvantaged elderly (Nogueira *et al.*, 2014).

Childhood maltreatment is widely associated with the risk of mental disease (Gilbert *et al.*, 2009; Taillieu *et al.*, 2016). Childhood maltreatment is associated with an increased incidence, a worse clinical course, and a poor response to the treatment of depression (Tunnard *et al.*, 2014; Wingo *et al.*, 2010).

The impact of childhood maltreatment seems to last a lifetime. In the elderly, it has a significant impact on physical and mental health (Draper *et al.*, 2008; Maschi *et al.*, 2012). Childhood maltreatment is extensively associated with the development of geriatric depression (Comijs *et al.*, 2013; Dannlowski *et al.*, 2012; Ege *et al.*, 2015; Kuhlman *et al.*, 2013; Raposo *et al.*, 2014). Thus, it is crucial to understand how childhood maltreatment impacts further development leading to pathology. It is also important to understand how multiple possible outcomes could be affected by personality traits.

Correspondence should be addressed to: Gabriel Behr Gomes Jardim, Instituto de Geriatria e Gerontologia - IGG, Pontifícia Universidade Católica do Rio Grande do Sul - PUCRS, Avenida Ipiranga 6681/703, Porto Alegre/RS, CEP: 90619-900, Brazil. Phone: +55 51 33536031. Email: gabrielbehr@gmail.com. Received 11 Oct 2018; revision requested 03 Nov 2018; revised version received 21 Dec 2018; accepted 27 Dec 2018. First published online 04 March 2019.

The big five personality traits (neuroticism, extroversion, agreeableness, conscientiousness, and openness) are associated with longevity and successful aging. Low scores of neuroticism and high scores of extroversion, openness, agreeableness, and conscientiousness are related to health characteristics of ageing with a high level of daily life activity and good cognitive outcomes (Baek *et al.*, 2016).

The big five personality traits are widely related to mental disease and childhood maltreatment. High levels of neuroticism and low levels of extraversion are associated with geriatric depression (Hayward, 2013; Koorevaar *et al.*, 2013; O'Shea *et al.*, 2017; Weber *et al.*, 2010). High levels of neuroticism are also associated with treatment resistance and cognitive symptomatology in those who have depression (Kendler *et al.*, 2006; Manning *et al.*, *in press*; Steffens *et al.*, 2017). Neuroticism is consistently associated with childhood maltreatment (Hengartner *et al.*, 2015; Mc Elroy and Hevey, 2014).

The literature on childhood maltreatment, geriatric depression, and personality traits suggests a mediating model with a temporal relationship between exposure to childhood maltreatment influencing the development of personality traits and increased risk of geriatric depression. These personality traits could be mediators that affect the risk of depression (Hovens *et al.*, 2016). This model is important because it could help develop treatments focused on personality traits that better treat geriatric depression.

The objective of this study is to test the big five personality traits as mediators of the relationship between childhood maltreatment and geriatric depression.

## Method

### Study sample

The Cerebral Aging Program (PENGE) is longitudinal cohort study that aims to examine mental health in a socioeconomically disadvantaged sample of elderly people. This sub-project is a cross-sectional study with a sample of persons 60 years old or older. This research was conducted from July 2015 to July 2016 in collaboration with the Family Health Strategy (FHS) program of Porto Alegre, Brazil.

General practitioners in family health facilities and health care workers in the PENGE program recruited the subjects. They were evaluated at the reference hospital. The inclusion criteria included being 60 years old or older. The exclusion criteria included the diagnosis of bipolar disorder; drug or alcohol abuse evaluated by the Mini International Neuropsychiatric Interview (MINI plus); history of neurological disease such as tumors, cerebral ischemia, and epilepsy; serious or incapacitating clinical disease that could

interfere with the interview such as cancer or visual defects; and those who had a cognitive inability to understand the questions (based on clinical judgment). Assessment and registry were made by board-certified psychiatrists and psychologists experienced in late-life neuropsychiatric disorders.

A total of 346 subjects were evaluated in the hospital, and 260 were included in the final sample after the inclusion and exclusion criteria were met and written informed consent collected. All subjects who had clinical needs were referred to specialized neurologic and psychiatric consultations in the same hospital.

The research protocol was fully approved by both the research ethics committees of PUCRS and the Public Health Secretary of the City of Porto Alegre (Brazil). All participants or their legal representatives gave informed written consent.

## Measurements

### Assessment of depression

Depression was measured with the Mini International Neuropsychiatric Interview 5.0 plus Portuguese version (MINI plus) (Amorim, 2000). The MINI plus is a gold-standard validated diagnostic tool. Its accuracy is similar to more complex psychiatric interviews in different settings (Lecrubier *et al.*, 1997). The major depression module is composed of questions based on the DSM IV criteria. MINI plus depression module was used as a latent variable composed of the MINI questions. The dependent variable used in the mediation model is composed of ten dichotomous questions (range 0-10).

### Assessment of childhood maltreatment

Childhood maltreatment was diagnosed with the shorter version of the Childhood Trauma Questionnaire (CTQ) (Bernstein *et al.*, 2003). It was translated into Brazilian Portuguese (Grassi-Oliveira *et al.*, 2006) and validated for community and clinical samples with good reliability and validity (Grassi-Oliveira *et al.*, 2014). The CTQ is composed of 28 questions that evaluate the subtype of maltreatment experience (emotional abuse, physical abuse, sexual abuse, emotional neglect, and physical neglect) as well as its intensity through a five-point Likert scale. The cut-off mild-high was used in accordance with the author's manual.

The five maltreatment subtypes include the following: Emotional abuse includes verbal assaults on a child's sense of worth or well-being or any humiliating behavior directed toward a child by an older person. Physical abuse includes bodily assaults on a child by an older person that confers a risk of or result

in injury. Sexual abuse is sexual conduct between a child and an adult or older person. Emotional neglect is the failure of an adult to meet the children's emotional and psychological needs. Physical neglect is the failure of adults to provide for a child's basic physical needs (food, shelter, clothing, safety, and health care).

### Assessment of personality traits

To assess personality factors, the NEO-Five Factor Inventory (NEO-FFI) (Costa and McCrae, 1995) was used. The NEO-FFI is a shortened version of the NEO Personality Inventory-Revised translated and validated to Brazilian Portuguese (Costa and McCrae, 2007). The 60-item questionnaire measures the main big five domains: neuroticism (emotionally unstable, susceptibility to psychological stress, maladjustment, and negative emotionality); extraversion (energetic, assertive); openness to experience (imaginative, independent-minded, intellectual curiosity); conscientiousness (responsible, dependable, orderly); and agreeableness (empathic, cooperative). The extent to which a statement applies to the participant is rated on a 5-point Likert scale ranging from "strongly disagree" to "strongly agree." The cut-off no-low was used in accordance with the author's manual.

### Assessments of sociodemographic variables

Age, gender, and education were used to describe the sample.

### Statistical analysis

Calculations used the software Statistical Package for the Social Sciences 20.0 SPSS®. Descriptive statistics were used to describe socioeconomic, personality traits, and history of childhood maltreatment.

We tested five independent mediation models using structural equation modeling (SEM), corrected normal theory method was performed using robust standard errors with weighted least squares estimation (WLS) and the corrected model test statistics using the Satorra-Bentler statistic (Kline, 2011). In each model, the independent variable was the total CTQ score. The dependent variable included questions from the MINI plus depression module. We used goodness-of-fit measures to evaluate how well the hypothesized model fit the observed data including  $\chi^2$ , Tucker Lewis index (TLI), comparative fit index (CFI), and root mean square error of approximation (RMSEA). A well-fitting model is seen when TLI or CFI is at least .90 and the RMSEA is .06 or lower. Statistical significance was considered to be  $P < 0.05$ .

## Results

### Descriptive Statistics

The sample included 260 participants with a mean age of 72.2 (SD 7.11) and a majority female (76.9%). The mean years of study (education level) were 4.72 (SD 3.75). Childhood maltreatment was common in this sample (physical neglect 50.4%; emotional neglect 99.2%; physical abuse 29.6%; emotional abuse 29.6%; and sexual abuse 13.5%). The personality traits presented high prevalence of high scores in this sample, except for openness. A high neuroticism score was seen in 69% of the sample ( $n = 180$ ) (Table 1). The five personality traits were correlated with childhood maltreatment: Extroversion, agreeableness, openness, and conscientiousness correlated positively; neuroticism correlated negatively.

### Mediation model

The independent model of mediation between childhood maltreatment and geriatric depression was tested via the big five personality characteristics. The full structural model showed acceptable fit:  $\chi^2(53) = 39.1$ ,  $P = 0.922$ ; RMSEA = 0.001; SRMR = 0.061; and CFI = 1.000. These are presented in Figure 1 and Table 2.

#### NEUROTICISM MEDIATION

Neuroticism has a strong total mediation effect. The CTQ score effect on the NEO-FFI neuroticism score was significant ( $\beta = 0.37$ , SE = 0.052,  $P < 0.001$ ), neuroticism is associated with MINI plus depression diagnosis ( $\beta = 0.66$ , SE = 0.050,  $P < 0.001$ ). The indirect effect of childhood maltreatment on depression was significant ( $\beta = 0.24$ , SE = 0.040,  $P < 0.001$ ). The direct effect of CTQ score on depression was negative and not significant ( $\beta = -0.06$ , SE = 0.061,  $P = 0.302$ ). The full model effect of CTQ score on depression was significant ( $\beta = 0.18$ , SE = 0.068,  $P = 0.009$ ).

#### EXTRAVERSION MEDIATION

Extraversion has a total mediation effect. The CTQ score effect on the NEO-FFI extraversion score was inverse and significant ( $\beta = -0.24$ , SE = 0.045,  $P < 0.001$ ); extraversion is associated with MINI plus depression diagnosis ( $\beta = -0.36$ , SE = 0.062,  $P < 0.001$ ). The indirect effect of childhood maltreatment on depression was significant ( $\beta = 0.09$ , SE = 0.022,  $P < 0.001$ ). The direct effect of CTQ score on depression was negative and not significant ( $\beta = 0.09$ , SE = 0.064,  $P = 0.138$ ). The full model effect of CTQ score on depression was significant ( $\beta = 0.18$ , SE = 0.068,  $P = 0.009$ ).

**Table 1.** Descriptive statistics ( $n = 260$ )

VARIABLE	MEAN (SD)(RANGE)	FREQUENCY, N (%)
<b>Socioeconomic Variables</b>		
Age (Years)	72.2(7,11)(60–95)	
Female Gender		200 (76.9%)
Education Level Years	4.72(3.75)(0–15)	
<b>Depression Diagnostic (MINI)</b>	2,55(3.32)(0–10)	83 (31.9%)
<b>Personality Characteristics (NEO-FFI)*</b>		
Neuroticism	37.6(9.25)(16–60)	180 (69.2%)
Extraversion	37.9(6.6)(21–56)	173 (66.5%)
Openness	34.9(3.9)(23–46)	67 (25.8%)
Agreeableness	40.8(3.82)(32–53)	216 (83.1%)
Conscientiousness	45.4(4.8)(29–59)	242 (93.1%)
<b>Childhood trauma (CTQ)**</b>		
CTQ Total	45.4(10.7)(31–105)	
Physical Neglect	8.5(3.6)(4–20)	131 (50.4%)
Emotional Neglect	15.7(1.9)(8–21)	258 (99.2%)
Physical Abuse	7.4(3.8)(4–25)	77 (29.6%)
Emotional Abuse	7.9(4.4)(4–25)	77 (29.6%)
Sexual Abuse	5.6(2.6)(4–25)	35 (13.5%)

Notes: Mini International Neuropsychiatric Interview (MINI); NEO-Five Factor Inventory (NEO-FFI); Childhood Trauma Questionnaire (CTQ).

\*Frequency: mild-high cut-off.

\*\*Frequency: no-low cut-off.

#### OPENNESS MEDIATION

There was no evidence for openness as a mediator. The CTQ score effect on NEO-FFI openness score was negative and significant ( $\beta = -0.14$ ,  $SE = 0.056$ ,  $P = 0.013$ ). Openness is associated with MINI plus depression diagnosis ( $\beta = -0.14$ ,  $SE = 0.063$ ,  $P = 0.022$ ). The indirect effect of childhood maltreatment on depression was not significant ( $\beta = 0.02$ ,  $SE = 0.012$ ,  $P = 0.091$ ). The direct effect of CTQ score on depression was significant ( $\beta = 0.16$ ,  $SE = 0.067$ ,  $P = 0.018$ ). The full model effect of CTQ score on depression was significant ( $\beta = 0.18$ ,  $SE = 0.068$ ,  $P = 0.009$ ).

#### AGREEABLENESS MEDIATION

Agreeableness has a partial mediation effect. The CTQ score effect on the NEO-FFI agreeableness score was negative and significant ( $\beta = -0.21$ ,  $SE = 0.058$ ,  $P < 0.001$ ). Neuroticism is associated with MINI plus depression diagnosis ( $\beta = -0.16$ ,  $SE = 0.068$ ,  $P = 0.017$ ). The indirect effect of childhood maltreatment on depression was significant ( $\beta = 0.03$ ,  $SE = 0.017$ ,  $P = 0.047$ ). The direct effect of CTQ score on depression was not significant ( $\beta = 0.15$ ,  $SE = 0.068$ ,  $P = 0.030$ ). The full model effect of CTQ score on depression was significant ( $\beta = 0.18$ ,  $SE = 0.068$ ,  $P = 0.009$ ).

#### CONSCIENTIOUSNESS MEDIATION

Conscientiousness has a partial mediation effect. The CTQ score effect on NEO-FFI conscientiousness score was negative and significant ( $\beta = -0.17$ ,


$SE = 0.048$ ,  $P < 0.001$ ). Conscientiousness is associated with the MINI plus depression diagnostic ( $\beta = -0.27$ ,  $SE = 0.062$ ,  $P < 0.001$ ). The indirect effect of childhood maltreatment on depression was significant ( $\beta = 0.05$ ,  $SE = 0.016$ ,  $P = 0.005$ ). The direct effect of CTQ score on depression was negative and not significant ( $\beta = 0.13$ ,  $SE = 0.066$ ,  $P = 0.041$ ). The full model effect of CTQ score on depression was significant ( $\beta = 0.18$ ,  $SE = 0.068$ ,  $P = 0.009$ ).

## Discussion

This study tested the hypothesis that personality traits mediate childhood maltreatment and depression in the elderly. The results demonstrated that neuroticism and extraversion are complete mediators; agreeableness and conscientiousness are partial mediators' and openness does not mediate the relationship.

### Socioeconomic profile of the population

This sample was mostly females who studied an average of four years. These individuals live in poor communities of Porto Alegre where violence, unemployment, and poor health are common. They tend to live in large families and deal with many stress factors daily. This environment can lead to increasing vulnerability at the edges of circle of life such as the elderly and children. This picture can be extended to a majority of the elderly population



**Figure 1.** The structural equation model of personality mediating childhood maltreatment and geriatric depression ( $n = 260$ ). Note: MINI questions, A1b Depressed mood; A2b Anhedonia; A3a Weight or appetite disturb; A3b Sleep disturb; A3c Psychomotor agitation or retardation; A3d Fatigue or loss of energy; A3e Worthlessness or guilt; A3f Concentration disturb; A3g Thoughts of death or suicidal ideation; A4 Daily functioning.  $*P \leq 0.05$ ,  $**P \leq 0.001$ .

who live in poor or developing countries—especially those in populous cities.

In these resource-scarce settings, elderly people often receive less health care, and children can be exposed to childhood maltreatment. The exposure to maltreatment decreases educational achievement and leads to low levels of employment and a significant loss of economic productivity. This can lead in turn to increased maltreatment (Boden *et al.*, 2007; Currie and Widom, 2010). This bidirectional relationship leads to a cycle of increasing vulnerability to mental disease.

### The mediation model of personality

Our results corroborate the widely studied association of childhood maltreatment and geriatric depression presented in the literature. In another study using the same population as this work, childhood abuse and neglect correlated with developing more severe forms of depression (Novelo *et al.*, 2018).

Although many depression treatments are available, they bring a number of difficulties to the patient such as a lack of efficacy and a poor side effect profile. Turning our attention to identifying

**Table 2.** Mediation analysis of personality characteristics (PC) on the relationship between childhood maltreatment (CM) score and depression (D)

MEDIATING VARIABLE	EFFECT OF CM ON PC (A)	EFFECT OF PC ON D (B)	INDIRECT EFFECT	DIRECT EFFECT OF CM ON D (C)	TOTAL EFFECT
<b>Neuroticism</b>	0,37**	0,66**	0.24**	-.06	0.18
<b>Extraversion</b>	-.24**	-.36**	0.09**	0.09	0.18
<b>Openness</b>	;-0.14*	-.14*	0.02	0.16	0.18
<b>Agreeableness</b>	-.21**	-.16*	0.03*	0.15*	0.18
<b>Conscientiousness</b>	-.17**	-.27**	0.05**	0.13*	0.18

\*P ≤ 0.05, \*\*P ≤ 0.001.

high-risk depression groups of individuals who were initially exposed to childhood maltreatment and have maladjusted personality traits could address new treatment options. (Beddington *et al.*, 2008; Kessler *et al.*, 2010; Ormel *et al.*, 2015)

We studied the big five personality traits because they are important constructs that seem to be robust, culturally invariable, and stable (Allik, 2005; Ormel *et al.*, 2015). They consistently correlate with childhood maltreatment (Hengartner *et al.*, 2015) and geriatric depression (Manning *et al.*, in press; O'Shea *et al.*, 2017; Steffens *et al.*, 2013).

Our results demonstrated that neuroticism and extraversion are complete mediators, agreeableness and conscientiousness are partial mediators, openness is not a mediator. In accordance with our findings, another study in adults showed that high levels of neuroticism and low levels of extraversion mediate the relationship between childhood maltreatment and a 4-year remission of depression and anxiety disorders (Hovens *et al.*, 2016). Similar studies showed that cognitive emotion dysregulation was an important mediator affecting depression and anxiety symptoms in patients with childhood maltreatment (Hopfinger *et al.*, 2016; Huh *et al.*, 2017).

Psychotherapy modalities focusing on high-risk individuals who experienced childhood maltreatment could also be studied. Probably, addressing as a psychotherapy objective the management of neuroticism's emotional instability, susceptibility to psychological stress, maladjustment, and negative emotionality could reduce the probability of experiencing geriatric depression. It could enhance extraversion, agreeableness, and conscientiousness.

Finally, enhancing life events that diminish the degree of maladaptive personality traits as good social support in adulthood, social relationship and work could be preventive strategies (Caspi *et al.*, 2005; Lüdtke *et al.*, 2011). For example, high resilience was related to improved physical well-being and metabolic biomarkers in schizophrenic individuals who suffered severe childhood trauma (Lee *et al.*, 2018).

### Strengths and limitations

The strength of this study is its focus on the mediation hypothesis of personality between childhood trauma and depression. To the best of our knowledge, there are only a few studies that evaluate this important subject. We are unaware of any studies that evaluate the psychopathology of the end of life with a focus on personality mediation factors. There are very few studies with socioeconomically disadvantaged elderly, but these persons represent the majority of global elderly population.

The study has some limitations. This cross-sectional design limits our conclusions on causality. Methodological limitations include retrospective reports of childhood trauma and personality characteristics measured at the same time. There is a tendency to recall bias and a tendency for the elderly to respond negatively to questions with aversive content. This can underestimate the results.

The prevalence of childhood maltreatment and personality traits might be higher in this population (MacDonald *et al.*, 2016). The measurement of personality traits at the same time as the childhood trauma refers to actual personality characteristics despite the already proven relative stability of these traits.

Data was collected in the hospital specific to the population. Some participants may not have been able to reach the facility. The transport structure available in the poor neighborhoods could also affect the mobility of some of the elderly subjects.

Finally, other mediator factors could influence the results including attachment characteristics, past history of psychopathology, or other important sociodemographic factors like education. Future studies that expand this mediation hypothesis to other variables are warranted.

### Conclusion

This study evaluates a model where personality characteristics mediate the relationship between childhood maltreatment and geriatric depression.

Mediation analysis showed that neuroticism and extraversion are complete mediators, agreeableness and conscientiousness are partial mediators, and openness is not a mediator. Treatments enhancing extraversion, agreeableness and conscientiousness, and diminishing neuroticism could reduce geriatric depression in high-risk elderly exposed to childhood trauma.

### Conflict of interest

None.

### Authors' roles

I. Gomes da Silva Filho, A. von Gunten, E. L. Nogueira, and A. Cataldo Neto designed the study, supervised the data collection, and supervised the writing of the manuscript. P. K. Ziegelmann and D. B. Bumaguin planned and performed the statistical analyses and supervised the writing of the manuscript. P. Engroff coordinated the data collection and the manuscript revision. G. B. G. Jardim collected data, analyzed the data, and wrote the manuscript.

### Acknowledgments

The authors would like to acknowledge the collaboration of professionals responsible for the Family Health Strategy from Porto Alegre city, Brazil.

This work was supported by CAPES (Coordenação de Aperfeiçoamento de Pessoal de Nível Superior – Brasil) scholarships and the PVE program (Programa Professor Visitante Especial) – finance code 001, PUCRS (Pontifícia Universidade Católica do Rio Grande do Sul).

### References

- Allik, J.** (2005). Personality dimensions across cultures. *Journal of Personality Disorders*, 19, 212–232.
- Amorim, P.** (2000). Mini International Neuropsychiatric Interview (MINI): validação de entrevista breve para diagnóstico de transtornos mentais Mini International Neuropsychiatric Interview (MINI): validation of a short structured diagnostic psychiatric interview. *Brazilian Journal of Psychiatry*, 22, 106–115.
- Baek, Y., Martin, P., Siegler, I. C., Davey, A. and Poon, L. W.** (2016). Personality traits and successful aging: findings from the Georgia Centenarian Study. *The International Journal of Aging and Human Development*, 83, 207–227. doi: [10.1177/0091415016652404](https://doi.org/10.1177/0091415016652404).
- Beddington, J. et al.** (2008). The mental wealth of nations. *Nature*, 455, 1057–1060. doi: [10.1038/4551057a](https://doi.org/10.1038/4551057a).
- Bernstein, D. P. et al.** (2003). Development and validation of a brief screening version of the Childhood Trauma Questionnaire. *Child Abuse and Neglect*, 27, 169–190. doi: [10.1016/S0145-2134\(02\)00541-0](https://doi.org/10.1016/S0145-2134(02)00541-0).
- Boden, J. M., Horwood, L. J. and Fergusson, D. M.** (2007). Exposure to childhood sexual and physical abuse and subsequent educational achievement outcomes. *Child Abuse and Neglect*, 31, 1101–1114. doi: [10.1016/j.chiabu.2007.03.022](https://doi.org/10.1016/j.chiabu.2007.03.022).
- Caspi, A., Roberts, B. W. and Shiner, R. L.** (2005). Personality development: stability and change. *Annual Review Psychology*, 56, 453–484. doi: [10.1146/annurev.psych.55.090902.141913](https://doi.org/10.1146/annurev.psych.55.090902.141913).
- Comijs, H. C., Van Exel, E., Van Der Mast, R. C., Paaauw, A., Oude Voshaar, R. and Stek, M. L.** (2013). Childhood abuse in late-life depression. *Journal of Affective Disorders*, 147, 241–246. doi: [10.1016/j.jad.2012.11.010](https://doi.org/10.1016/j.jad.2012.11.010).
- Conwell, Y., Duberstein, P. R. and Caine, E. D.** (2002). Risk factors for suicide in later life. *Biological Psychiatry*, 52, 193–204. doi: [10.1016/S0006-3223\(02\)01347-1](https://doi.org/10.1016/S0006-3223(02)01347-1).
- Costa, P. and McCrae, R. R.** (1995). Inventory domains and facets: hierarchical personality assessment using the revised NEO personality inventory. *Journal of Personality Assessment*, 64, 21–50. doi: [10.1207/s15327752jpa6401](https://doi.org/10.1207/s15327752jpa6401).
- Currie, J. and Widom, C. S.** (2010). Long-term consequences of child abuse and neglect on adult economic well-being. *Child Maltreatment*, 14, 384–399. doi: [10.1080/10810730902873927](https://doi.org/10.1080/10810730902873927). Testing.
- Dannowski, U. et al.** (2012). Limbic scars: long-term consequences of childhood maltreatment revealed by functional and structural magnetic resonance imaging. *Biological Psychiatry*, 71, 286–293. doi: [10.1016/j.biopsych.2011.10.021](https://doi.org/10.1016/j.biopsych.2011.10.021).
- Draper, B. et al.** (2008). Long-term effects of childhood abuse on the quality of life and health of older people: results from the depression and early prevention of suicide in general practice project. *Journal of the American Geriatrics Society*, 56, 262–271. doi: [10.1111/j.1532-5415.2007.01537.x](https://doi.org/10.1111/j.1532-5415.2007.01537.x).
- Dube, S. R., Felitti, V. J., Dong, M., Giles, W. H. and Anda, R. F.** (2003). The impact of adverse childhood experiences on health problems: evidence from four birth cohorts dating back to 1900. *Preventive Medicine*, 37, 268–277. doi: [10.1016/S0091-7435\(03\)00123-3](https://doi.org/10.1016/S0091-7435(03)00123-3).
- Ege, M. A., Messias, E., Thapa, P. B. and Krain, L. P.** (2015). Adverse childhood experiences and geriatric depression: results from the 2010 BRFSS. *The American Journal of Geriatric Psychiatry*, 23, 110–114. doi: [10.1016/j.jagp.2014.08.014](https://doi.org/10.1016/j.jagp.2014.08.014).
- Gilbert, R., Widom, C. S., Browne, K., Fergusson, D., Webb, E. and Janson, S.** (2009). Burden and consequences of child maltreatment in high-income countries. *The Lancet*, 373, 68–81. doi: [10.1016/S0140-6736\(08\)61706-7](https://doi.org/10.1016/S0140-6736(08)61706-7).
- Grassi-Oliveira, R. et al.** (2014). Childhood Trauma Questionnaire (CTQ) in Brazilian samples of different age groups: findings from confirmatory factor analysis. *PLoS ONE*, 9, E87118. doi: [10.1371/journal.pone.0087118](https://doi.org/10.1371/journal.pone.0087118).
- Grassi-Oliveira, R., Stein, L. M. and Pezzi, J. C.** (2006). Tradução e validação de conteúdo da versão em português do Childhood Trauma Questionnaire. *Revista de*

- Saude Publica*, 40, 249–255. doi: [10.1590/S0034-89102006000200010](https://doi.org/10.1590/S0034-89102006000200010).
- Hayward, R. D.** (2013). Association of NEO personality. *American Journal of Geriatric Psychiatry*, 21, 88–96. doi: [10.1016/j.jagp.2012.11.012](https://doi.org/10.1016/j.jagp.2012.11.012).
- Hengartner, M. P., Cohen, L. J., Rodgers, S., Müller, M., Rössler, W. and Ajdacic-Gross, V.** (2015). Association between childhood maltreatment and normal adult personality traits: exploration of an understudied field. *Journal of Personality Disorders*, 29, 1–14. doi: [10.1521/pedi\\_2014\\_28\\_143](https://doi.org/10.1521/pedi_2014_28_143).
- Hopfinger, L., Berking, M., Bockting, C. L. H. and Ebert, D. D.** (2016). Emotion regulation mediates the effect of childhood trauma on depression. *Journal of Affective Disorders*, 198, 189–197. doi: [10.1016/j.jad.2016.03.050](https://doi.org/10.1016/j.jad.2016.03.050).
- Hovens, J. G. F. M., Giltay, E. J., Van Hemert, A. M. and Penninx, B. W. J. H.** (2016). Childhood maltreatment and the course of depressive and anxiety disorders: the contribution of personality characteristics. *Depression and Anxiety*, 33, 27–34. doi: [10.1002/da.22429](https://doi.org/10.1002/da.22429).
- Huh, H. J., Kim, K. H., Lee, H. K. and Chae, J. H.** (2017). The relationship between childhood trauma and the severity of adulthood depression and anxiety symptoms in a clinical sample: the mediating role of cognitive emotion regulation strategies. *Journal of Affective Disorders*, 213, 44–50. doi: [10.1016/j.jad.2017.02.009](https://doi.org/10.1016/j.jad.2017.02.009).
- Kendler, K. S., Gatz, M., Gardner, C. O. and Pedersen, N. L.** (2006). Personality and major depression. *Archives of General Psychiatry*, 63, 1113. doi: [10.1001/archpsyc.63.10.1113](https://doi.org/10.1001/archpsyc.63.10.1113).
- Kessler, R. C. et al.** (2010). Age differences in the prevalence and co-morbidity of DSM-IV major depressive episodes: results from the WHO world mental health survey initiative. *Depression and Anxiety*, 27, 351–364. doi: [10.1002/da.20634](https://doi.org/10.1002/da.20634).
- Kline, R. B.** (2011). *Principles and Practice of Structural Equation Modeling* (3rd edn). New York, NY: Guilford Press.
- Koorevaar, A. M. L. et al.** (2013). Big five personality and depression diagnosis, severity and age of onset in older adults. *Journal of Affective Disorders*, 151, 178–185. doi: [10.1016/j.jad.2013.05.075](https://doi.org/10.1016/j.jad.2013.05.075).
- Kuhlman, K. R., Maercker, A., Bachem, R., Simmen, K. and Burri, A.** (2013). Developmental and contextual factors in the role of severe childhood trauma in geriatric depression: the sample case of former indentured child laborers. *Child Abuse and Neglect*, 37, 969–978. doi: [10.1016/j.chiabu.2013.04.013](https://doi.org/10.1016/j.chiabu.2013.04.013).
- Lecrubier, Y. et al.** (1997). The Mini International Neuropsychiatric Interview (MINI). A short diagnostic structured interview: reliability and validity according to the CIDI. *European Psychiatry*, 12, 224–231. doi: [10.1016/S0924-9338\(97\)83296-8](https://doi.org/10.1016/S0924-9338(97)83296-8).
- Lee, E. E., Martin, A. S., Tu, X., Palmer, B. W. and Jeste, D. V.** (2018). Childhood adversity and schizophrenia: the protective role of resilience in mental and physical health and metabolic markers. *Journal of Clinical Psychiatry*, 79, 1–9.
- Lüdtke, O., Roberts, B. W. and Nagy, G.** (2011). A random walk down university avenue: life paths, life events, and personality trait change at the transition to university life. *Journal of Personality and Social Psychology*, 101, 620–637. doi: [10.1037/a0023743.A](https://doi.org/10.1037/a0023743.A).
- MacDonald, K. et al.** (2016). Minimization of childhood maltreatment is common and consequential: results from a large, multinational sample using the Childhood Trauma Questionnaire. *PLoS ONE*, 11, 1–16. doi: [10.1371/journal.pone.0146058](https://doi.org/10.1371/journal.pone.0146058).
- Manning, K. J., Ph, D., Chan, G., Ph, D. and Steffens, D. C.** (2016). Neuroticism traits selectively impact long-term illness course and cognitive decline in late-life depression. *The American Journal of Geriatric Psychiatry*, 25, 1–10. doi: [10.1016/j.jagp.2016.10.006](https://doi.org/10.1016/j.jagp.2016.10.006).
- Maschi, T., Baer, J., Morrissey, M. B. and Moreno, C.** (2012). The aftermath of childhood trauma on late life mental and physical health: a review of the literature. *Traumatology*, 19, 1–6. doi: [10.1177/1534765612437377](https://doi.org/10.1177/1534765612437377).
- McElroy, S. and Hevey, D.** (2014). Relationship between adverse early experiences, stressors, psychosocial resources and wellbeing. *Child Abuse and Neglect*, 38, 65–75. doi: [10.1016/j.chiabu.2013.07.017](https://doi.org/10.1016/j.chiabu.2013.07.017).
- Nogueira, E. L., Rubin, L. L., Giacobbo, S. de, S., Gomes, I. and Neto, A. C.** (2014). Screening for depressive symptoms in older adults in the Family Health Strategy, Porto Alegre, Brazil. *Revista de Saude Publica*, 48, 368–377. doi: [10.1590/S0034-8910.2014048004660](https://doi.org/10.1590/S0034-8910.2014048004660).
- Novelo, M. et al.** (2018). Effects of childhood multiple maltreatment experiences on depression of socioeconomic disadvantaged elderly in Brazil. *Child Abuse and Neglect*, 79, 350–357. doi: [10.1016/j.chiabu.2018.02.013](https://doi.org/10.1016/j.chiabu.2018.02.013).
- O'Shea, D. M., Dotson, V. M. and Fieo, R. A.** (2017). Aging perceptions and self-efficacy mediate the association between personality traits and depressive symptoms in older adults. *International Journal of Geriatric Psychiatry*, 32, 1217–1225. doi: [10.1002/gps.4584](https://doi.org/10.1002/gps.4584).
- Ormel, J. et al.** (2015). Neuroticism and common mental disorders: meaning and utility of a complex relationship. *Clinical Psychology Review*, 33, 686–697. doi: [10.1016/j.cpr.2013.04.003](https://doi.org/10.1016/j.cpr.2013.04.003).
- Penninx, B. W., Geerlings, S. W., Deeg, D. J., van Eijk, J. T., van Tilburg, W. and Beekman, A. T.** (1999). Minor and major depression and the risk of death in older persons. *Archives of General Psychiatry*, 56, 889–895. doi: [10.1001/archpsyc.56.10.889](https://doi.org/10.1001/archpsyc.56.10.889).
- Raposo, S. M., Mackenzie, C. S., Henriksen, C. A. and Afifi, T. O.** (2014). Time does not heal all wounds: older adults who experienced childhood adversities have higher odds of mood, anxiety, and personality disorders. *American Journal of Geriatric Psychiatry*, 22, 1241–1250. doi: [10.1016/j.jagp.2013.04.009](https://doi.org/10.1016/j.jagp.2013.04.009).
- Sachs-Ericsson, N. J., Rushing, N. C., Stanley, I. H. and Sheffler, J.** (2016). In my end is my beginning: developmental trajectories of adverse childhood experiences to late-life suicide. *Aging & Mental Health*, 20, 139–165. doi: [10.1080/13607863.2015.1063107](https://doi.org/10.1080/13607863.2015.1063107).
- Steffens, D. C., McQuoid, D. R., Smoski, M. J. and Potter, G. G.** (2013). Clinical outcomes of older depressed patients with and without comorbid neuroticism. *International Psychogeriatrics*, 25, 1985–1990. doi: [10.1017/S1041610213001324](https://doi.org/10.1017/S1041610213001324).
- Steffens, D. C., Wu, R., Grady, J. J. and Manning, K. J.** (2017). Presence of neuroticism and antidepressant



remission rates in late-life depression: results from the Neurobiology of Late-Life Depression (NBOLD) study. *International Psychogeriatrics*, 30, 1069–1074. doi: [10.1017/S1041610217002551](https://doi.org/10.1017/S1041610217002551).

**Taillieu, T. L., Brownridge, D. A., Sareen, J. and**

**Affi, T. O.** (2016). Child abuse & neglect childhood emotional maltreatment and mental disorders: results from a nationally representative adult sample from the United States. *Child Abuse & Neglect*, 59, 1–12. doi: [10.1016/j.chiabu.2016.07.005](https://doi.org/10.1016/j.chiabu.2016.07.005).

**Tunnard, C., Rane, L. J., Wooderson, S. C.,**

**Markopoulou, K., Poomb, L., Fekadu, A., Juruena, M.,**

**Cleare, A. J.** (2014). The impact of childhood adversity on suicidality and clinical course in treatment-resistant depression. *Journal of Affective Disorders*, 152–154, 122–130. doi: [10.1016/j.jad.2013.06.037](https://doi.org/10.1016/j.jad.2013.06.037).

**Weber, K. et al.** (2010). Volumetric MRI changes, cognition and personality traits in old age depression. *Journal of Affective Disorders*, 124, 275–282. doi: [10.1016/j.jad.2009.11.016](https://doi.org/10.1016/j.jad.2009.11.016).

**Wingo, A. P., Wrenn, G., Pelletier, T., Gutman, A. R., Bradley, B. and Ressler, K. J.** (2010). Moderating effects of resilience on depression in individuals with a history of childhood abuse or trauma exposure. *Journal of Affective Disorders*, 126, 411–414. doi: [10.1016/j.jad.2010.04.009](https://doi.org/10.1016/j.jad.2010.04.009).