



Three-dimensional virtual planning for nodule resection in solid organs: A systematic review and meta-analysis

Matheus Zanon^{a,b,*}, Stephan Altmayer^{b,c}, Guilherme Watte^{a,b,c}, Gabriel Sartori Pacini^b, Tan-Lucien Mohammed^d, Edson Marchiori^e, Darcy Ribeiro Pinto Filho^f, Bruno Hochhegger^{a,b,c}

^a Graduate Program in Pathology, Federal University of Health Sciences of Porto Alegre - R. Sarmento Leite, 245, Porto Alegre, 90050170, Brazil

^b Medical Imaging Research Lab, LABIMED, Department of Radiology, Pavilhão Pereira Filho Hospital, Irmandade Santa Casa de Misericórdia de Porto Alegre - Av. Independência, 75, Porto Alegre, 90020160, Brazil

^c Postgraduate Program in Medicine and Health Sciences, Pontifícia Universidade Católica do Rio Grande do Sul, Av. Ipiranga, 6690, Porto Alegre, 90619900, Brazil

^d Department of Radiology, College of Medicine, University of Florida, 1600 SW Archer Rd m509, Gainesville, FL, 32610, United States

^e Department of Radiology, Federal University of Rio de Janeiro - Av. Carlos Chagas Filho, 373, Rio de Janeiro, 21941902, Brazil

^f Department of Thoracic Surgery, University of Caxias do Sul, R. Francisco Getúlio Vargas, 1130, Caxias do Sul, 95070561, Brazil

ARTICLE INFO

Keywords:

Three-dimensional
Image-guided surgery
Lung
Liver
Kidney

ABSTRACT

Objectives: To systematically review the effects of 3D-imaging virtual planning for nodule resection in the following solid organs: lung, liver, and kidney.

Methods: MEDLINE, EMBASE, and Cochrane Library were searched through September 31, 2020 to include randomized and non-randomized controlled studies that compared outcomes of surgical resection of lung, liver, or kidney nodule resection with and without 3D virtual planning with computed tomography. From each article, the mean operation time (OT), mean estimated blood loss (EBL), mean postoperative hospital stay (POHS), and the number of postoperative events (POE) were extracted. The effect size (ES) of 3D virtual planning vs. non-3D planning was extracted from each study to calculate the pooled measurements for continuous variables (OT, EBL, POHS). Data were pooled using a random-effects model.

Results: The literature search yielded 2397 studies and 10 met the inclusion criteria with a total of 897 patients. There was a significant difference in OT between groups with a moderate ES favoring the 3D group (ES, -0.56; 95%CI: 0.91, -0.29; $I^2 = 83.1\%$; $p < .001$). Regarding EBL, there was a significant difference between 3D and non-3D with a small ES favoring IGS (ES, -0.18; 95%CI: 0.33, -0.02; $I^2 = 22.5\%$; $p = .0236$). There was no difference between the 3D and non-3D groups for both POHS (POHS ES, -0.15; 95%CI: 0.39, 0.10; $I^2 = 37.0\%$; $p = .174$) and POE (POE odds ratio (OR), 0.80; 95%CI: 0.54, 1.19; $I^2 = 0.0\%$; $p = .0973$).

Conclusions: 3D-imaging planning for surgical resection of lung, kidney, and liver nodules could reduce OT and EBL with no effects on immediate POHS and POE. Improvements in these perioperative variables could improve medium and long-term postoperative clinical outcomes.

1. Introduction

With the development of new post-processing technologies, there has been an increase in the use of image-guided surgery (IGS) systems to assist surgeons during challenging operations [1,2]. Several preoperative and intraoperative applications of IGS systems are currently available. The most used IGS systems are those that provide preoperative

information support, such as segmentation of risk structures or targets by three-dimensional (3D) virtual guidance. IGS systems can also offer intraoperative information support, such as distance visualization, proximity warnings when close to important structures, and augmentation of risk structures or targets [1,2].

IGS systems could improve intraoperative orientation, identification, and location of anatomical structures and their variations, decreasing

* Corresponding author. Department of Diagnostic Methods, Federal University of Health Sciences of Porto Alegre - R. Sarmento Leite, 245, Porto Alegre, 90050-170, Brazil.

E-mail addresses: mhgzanon@hotmail.com, mzanon200@caledonian.ac.uk (M. Zanon), stephanaltmayer@gmail.com (S. Altmayer), g.watte@gmail.com (G. Watte), gabrielsartorimed@gmail.com (G.S. Pacini), mohtan@radiology.ufl.edu (T.-L. Mohammed), edmarchiori@gmail.com (E. Marchiori), darcyryp@terra.com.br (D.R. Pinto Filho), brunoho@ufcspa.edu.br (B. Hochhegger).

<https://doi.org/10.1016/j.suronc.2021.101598>

Received 22 October 2020; Received in revised form 8 April 2021; Accepted 26 April 2021

Available online 30 April 2021

0960-7404/© 2021 Published by Elsevier Ltd.

surgeons' workload and contributing to performance enhancement [1, 3–7]. On the other hand, adverse side effects could also arise from the use of IGS systems, such as increased perceived time, pressure and mental demands, errors due to overreliance on the computer assistance, and possible interference on the development of surgical skills [2].

Clinical outcomes, such as patient safety, operation time (OT), intraoperative estimated blood loss (EBL), postoperative hospital stay (POHS), and prevalence of postoperative events (POE) have been reported to improve with the use of IGS systems [1,8–11]. Previous systematic reviews and meta-analyses have reported positive effects of IGS on clinical outcomes of maxillofacial surgery, neurosurgery, and spinal surgery [3–5]. Until now, there are only a few state-of-the-art reviews about the use of 3D-guidance for operations of the lung [12–14], liver [15–17], and kidney [18–21], but no systematic reviews or meta-analyses on the effect of IGS on these solid organs were performed. This meta-analysis aimed to assess the effects of 3D-imaging virtual planning (3DVP) for surgical resection of lung, liver, and kidney nodules.

2. Methods

2.1. Search strategy

This study was reported following Enhancing the Quality and Transparency of Health Research Reporting Guidelines, including the Preferred Reporting Items for Systematic Reviews and Meta-analysis Of Observational Studies in Epidemiology guidelines. We searched all available literature published in the PubMed-MEDLINE, EMBASE, and Cochrane databases through September 31, 2020. The databases were comprehensively searched using the equivalent terms included in [Appendix E1](#).

2.2. Inclusion and exclusion criteria

Studies were eligible for inclusion if the following criteria were present [1]: comparison of surgical outcomes of lung, liver, or kidney nodule resection surgery with and without image guidance [2]; report of the 3DVP effects on at least one of the following clinical outcomes: operation time, intraoperative bleeding, postoperative hospital stay, or postoperative events [3]; use of preoperative or intraoperative 3D reconstruction from computed tomography (CT) or magnetic resonance images; and [4] design of the study as randomized and non-randomized controlled trials.

Exclusion criteria were the following [1]: studies that used image guidance for biopsy, radiotherapy, or robotic-assisted surgery [2]; studies that used other types of IGS, not the 3D reconstruction, such as augmented reality [3]; case reports, letters to editor, reviews, or meta-analysis [4]; studies not published in English [5]; studies with animals, corpses, or phantoms.

2.3. Study quality assessment

Two reviewers assessed the quality of all eligible studies with the methodological index for non-randomized studies (MINORS) [22]. This tool is composed of 12 items that assess the methodological quality of non-randomized surgical studies. Each of these items is scored from 0 to 2, where 0 indicates that the issue was not reported in the evaluated study, 1 corresponding to items reported inadequately, and 2 correspond to items reported adequately [22].

2.4. Data extraction

Three reviewers independently reviewed all included articles to collect all the primary data (e.g., study design, country of recruitment, type of surgery, nodule site). From each article, the mean operation time (OT), mean estimated blood loss (EBL), mean postoperative hospital stay

(POHS), and the number of postoperative events (POE) were extracted. Any disagreements were resolved by consensus.

2.5. Statistical analysis

The effect size (ES) (standardized mean difference, SMD) of 3DVP versus non-3DVP was extracted from each study to calculate the pooled measurements for continuous variables (OT, EBL, POHS). The magnitude of effect for SMD was considered as “small” if equal to 0.2, “medium” if 0.5, and large if 0.8 [23]. The odds ratio (OR) were extracted from each study to combine the pooled results regarding the number of POE. In studies where the standard deviation was not reported, it was estimated based on sample size, median and interquartile range [24]. Data were pooled using a random-effects model.

Heterogeneity between studies was tested with the Q test [25]. The I^2 index was used to quantify the extent of heterogeneity. Publication bias was estimated using the funnel plot and the Egger's and Begg's tests [26, 27]. Sensitivity analyses that excluded each of the individual articles were conducted to evaluate whether any specific study significantly influenced the overall pooled results. All P-values less than 0.05 were considered statistically significant. All statistical analyses were performed using Stata version 15.0 (StataCorp LP, College Station, Texas, USA).

3. Results

3.1. Study characteristics


The initial search yielded 2397 studies, from which 86 were reviewed, and 10 met the inclusion criteria [8–11,28–33] ([Fig. 1](#)). The median MINORS score was 17 (interquartile range (IQR), 15.25–17.25). Only one study had a MINORS score lower than 15 (MINORS score = 5) [28].

A total of 897 patients were included, of which 469 (52.3%) subjects had undergone 3DVP, and 428 (47.7%) were non-3DVP controls ([eTable 1](#)). Lung was the main surgical site in two studies that performed video-assisted thoracoscopic (VATS) segmentectomy, lobectomy, and bilobectomy [8,28]. Three studies approached liver nodule resection using 3DVP, performing hemihepatectomy, extended hepatectomy, segmentectomy, sectionectomy, or partial resection [9,29,30]. Five studies used image-guided kidney surgery using laparoscopic partial nephrectomy (LPN), robot-assisted partial nephrectomy, or minimum incision endoscopic nephrectomy [10,11,31–33].

3.2. Operation time

All studies were included in the pooled OT analysis [8–11,28–33] ([Fig. 2](#)). Six studies described a significant shorter OT for IGS [8–11,29,30]. In the remaining, there was no significant difference in the OT between IGS and non-IGS [28,31–33]. Pooled analysis revealed a significant difference on OT between the groups with a moderate ES favoring the IGS group and a significantly high heterogeneity between studies (SMD, -0.56 ; 95%CI: $-0.91, -0.22$; $I^2 = 83.1\%$; $p < .001$). Visual analysis of the forest plot ([Fig. 2](#)) revealed that the article by Xue et al. was the greater contributor to the high heterogeneity between studies. Therefore, we performed a sensitivity analysis with removal of such study, which resulted in a lower but still significant heterogeneity ($I^2 = 52.8\%$; $p = .031$), and a slightly lower effect size (SMD, -0.36 ; 95%CI: $-0.58, -0.15$) ([Fig. 3](#)).

Nakayama et al. also performed a subgroup analysis of the OT and found that IGS impact was more significant for patients with repeated hepatectomy, as the 3D group presented a 130 min shorter median OT compared to the group without 3D ($p = .03$) [9]. Among the types of hepatic resections, segmentectomy was the only that presented a statistically significant shorter OT for the 3D group (MD, -43.5 min) ($p = .03$) [9].



* Note. - The same study could be excluded for multiple reasons.

Fig. 1. Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) flow diagram.




Fig. 2. Forest plot of standard mean difference for operation time.




Fig. 3. Sensitivity analysis of standard mean difference for operation time.

Although Wang et al. did not find a significant difference in the mean OT between 3D-guided and non-3D-guided LPN, the authors found a significant shorter OT (MD = -28.1 min; p = .018) for patients with medium-high complexity renal lesions that had a 3D virtual-surgery planning (i.e., R.E.N.A.L. score ≥ 8) [31]. Likewise, when controlling for case complexity and other covariates, Shirk et al. also found that patients whose surgical planning involved 3-D VR models showed differences in OT (OR, 1.00; 95%CI, 0.37–2.70; estimated OR, 2.47).

3.3. Estimated blood loss

All studies were included in the pooled EBL analysis [8–11,28–33]. Only two studies found a significantly lower EBL for 3DVP [10,11]. There was a significant difference between 3DVP and non-3DVP with a

small ES favoring IGS and a non-significant heterogeneity between studies (SMD, -0.18; 95%CI: -0.33, -0.02; $I^2 = 22.5\%$; p = .236) (Fig. 4).

Wang et al. reported a mean EBL of 148.1 (range, 85–290) mL for 3D-guided LPN vs. a mean EBL of 176.1 (range, 80–225) mL for non-3D-guided LPN (p < .001) [11]. However, in our pooled analysis, the SMD between the groups in this study was not significant (SMD, -0.56; 95%CI: -1.25, 0.13).

Although Shirk et al. did not find significant differences in EBL between the control and intervention groups, patients without 3D VR-assisted surgical planning were more likely to have EBL greater than 200 mL (OR, 1.98; 95%CI, 1.04–3.78) [33]. In addition, when controlling for case complexity and other covariates, the non-IGS group was more likely to have higher EBL (OR, 1.98; 95%CI, 1.04–3.78; estimated




Fig. 4. Forest plot of standard mean difference for estimated blood loss.

OR, 4.56) [33].

3.4. Postoperative hospital stay and postoperative events

Five studies reported the POHS [8,9,11,28,30], and six studies reported POE [8,9,29–32]. None of the studies reported a statistically significant difference in the length of POHS or the number of POE between 3DVP and non-3DVP. The pooled analyses failed to demonstrate any difference between the 3DVP and non-3DVP groups for both POHS (SMD, -0.15; 95%CI: -0.39, 0.10; I² = 37.0%; p = .174) (Fig. 5) and POE (OR, 0.80; 95%CI: 0.54, 1.19; I² = 0.0%; p = .0973) (Fig. 5).

For lung surgery, PO complications occurred at similar rates for 3D-guided (n = 6; 17%) and non-3D-guided VATS-segmentectomy (n = 5; 16%) in the study by Xue et al. [8]. These included atrial fibrillation, air leakage, pneumonia, and atelectasis. Xue et al. also reported a similar chest tube duration (3D-VATS-segmentectomy, 4.1 ± 1.8 days; vs. non-3D-VATS-segmentectomy, 4.1 ± 2.2 days; p = .93) [8].

Nakayama et al. only defined POE as complications of grade IIIa or higher by the Clavien-Dindo classification [9,34]. Although Fang et al. did not find any significant difference for POE between the groups,

non-3D-guided liver resections had significantly more Clavien-Dindo complications of grades III and IV (n = 8; 14.3%; vs. n = 2; 3.3%; p = .048) [30]. Patients that underwent 3D-guided liver resection also had less PO ascites (n = 2; 3.3%; vs. n = 8; 14.3%; p = .048), lower serum total bilirubin (23.2 ± 16.1 g/L; vs. 31.1 ± 24.1 g/L; p = .032), and higher serum albumin (29.3 ± 5.2 g/L; vs 27.8 ± 7.9 g/L; p = .0330) [30].

Two studies on renal nodule resection reported a lower rate of opening of the collecting system and urinary leakage in the 3D-guided nephrectomy groups [31,32]. Wang et al. reported significantly more cases of urinary leakage for patients with medium-high complexity renal lesions (i.e., R.E.N.A.L. score ≥8) that underwent non-3D-guided nephrectomy (4 vs. 0; p = .033) [31]. None of the studies reported significant differences between the IGS and non-IGS groups on renal function, such as a significant increase in serum creatinine level or decrease in ipsilateral glomerular filtration rate postoperatively [31,32]. Although Shirk et al. did not report median POHS for the control and intervention groups, the authors found that patients without 3D VR-assisted surgical planning were more likely to have a length of hospital stay longer than two days (OR, 2.86; 95%CI, 1.59–5.14) [33].




Fig. 5. Forest plots of (A) standard mean difference for postoperative hospital stay and (B) odds ratio for postoperative events.

Further, when controlling for case complexity and other covariates, patients whose surgical planning involved 3-D VR models showed differences in clamp time (OR, 1.60; 95%CI, 0.79–3.23; estimated OR, 11.22), and length of hospital stay (OR, 2.86; 95%CI, 1.59–5.14; estimated OR, 5.43) [33].

4. Discussion

In this meta-analysis, we found that IGS resulted in significantly lower OT and EBL but had no significant impact on POHS and POE. These results demonstrate that 3D-imaging virtual planning could improve some perioperative surgical variables. Although this meta-analysis could not find significant differences in the immediate postoperative clinical outcomes (POHS and POE), improvements in perioperative variables such as OT and EBL could influence medium and long-term clinical outcomes [35–42].

Previous data have shown that prolonged OT was associated with increased risk of the surgical site. In a previous meta-analysis, every additional 10 min, 15 min, 30 min, and 60 min of surgery increased the likelihood of surgical site infection in 5%, 13%, 17%, and 37%, respectively [35]. In another study including over 76,000 patients who underwent laparoscopic procedures, increasing operation time was independently linked with increased odds of complication in several elective procedures, such as colectomy, cholecystectomy, and gastric bypass (⁴²).

Also, previous studies have reported worse outcomes associated with higher blood losses during surgery, especially in the kidney [43], liver [44–47], and lung [48–50]. In one series evaluating partial nephrectomy, operative blood loss higher than 250 mL and 1000 mL were associated with 60% and 1150% increased risk of hemorrhagic complications, respectively (⁴³). For laparoscopic liver resection, EBL >250 mL has been described to correlate with higher conversion rates to open surgery and overall complications, including liver and kidney failure and postoperative mortality (⁴⁴). In another series including resection of hepatocellular carcinoma, higher blood losses were associated with tumor recurrence and survival (⁴⁷). In our study, we found a small but significant effect size on the use of 3DVP towards minimizing bleeding during surgery. Although this effect was small, techniques that minimize blood losses may benefit nodule resection surgery, reducing complications and postoperative mortality. As 3DVP allows a throughout preoperative evaluation of blood supply and anatomy variations, large blood losses could be prevented, enhancing patient safety.

The use of IGS systems was linked to a reduction of 52% in the number of significant complications and 34% of total complications in a previous meta-analysis on image-guided endoscopic sinus surgery [3]. In another meta-analysis of pedicle screw insertion in spine surgery, the use of IGS systems was associated with a reduction of 61% in the risk of pedicle perforation [51]. However, data for other types of surgery are scarce. To our knowledge, this is the first meta-analysis evaluating only solid organs.

Some concerns have been raised related with the use of IGS systems, as workload increase and impairment on the learning of new surgical skills. For image-guided endoscopic sinus surgery, the previous meta-analysis could not find any significant difference between image-guided and standard surgeries, even when the included studies objectively assessed cardiovascular and endocrine indicators of workload and stress [3]. Also, 3D reconstructions have been shown to impact positively on the surgical education in some papers [52,53]. In our meta-analysis, we did not assess the 3DVP impact on surgeons, but future studies should try to include such analysis.

Our study has some limitations. First, our pooled analysis demonstrated significantly high heterogeneity for the OT ($I^2 = 83.1\%$; $p = .000$) that was reduced in a complementary analysis without the study by Xue et al. ($I^2 = 52.8\%$; $p = .031$) without the loss of the favoring effect towards IGS. Such findings could be attributed to several factors, including the comparison of different organ surgeries, types of surgery,

sample sizes, surgeons' experience, software used for reconstruction, and different levels of surgery complexity. However, secondary subgroup analysis to try to identify the sources of heterogeneity, including those mentioned above, could not be performed due to the limited number of studies included in the meta-analysis. However, by using standardized mean differences, we tried to minimize the influence of some of these variables in our analysis. Second, we only included three solid organs that are among the most common sites of resection surgery of nodular lesions. Future studies using other organs could help to validate the benefits of IGS systems further. Third, there were limitations inherent to any meta-analyses, such as selection bias, publication bias, missing information from studies. Finally, most of the included studies were not prospective randomized clinical trials comparing IGS vs. non-IGS.

In summary, our data demonstrated that 3D-imaging virtual planning for resection of lung, kidney, and liver nodules could reduce operation time and estimated blood loss with no effects on immediate postoperative hospital stay and postoperative events.

Funding

This work did not receive any funding.

Availability of data and material

Data will be provided online once the article after the article is accepted.

Ethical approval

All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards. For this type of study formal consent is not required.

Informed consent

This article does not contain patient data.

Contributions

Matheus Zanon: Conceptualization; Data curation; Formal analysis; Investigation; Methodology; Project administration; Resources; Supervision; Validation; Visualization; Roles/Writing - original draft; Writing - review & editing.

Stephan Altmayer: Formal analysis; Investigation; Methodology; Validation; Visualization; Roles/Writing - original draft.

Guilherme Watte: Formal analysis; Investigation; Methodology; Validation; Visualization; Software; Roles/Writing - original draft.

Gabriel Sartori Pacini: Formal analysis; Investigation; Methodology; Validation; Visualization; Roles/Writing - original draft.

Tan-Lucien Mohammed: Conceptualization; Data curation; Formal analysis; Investigation; Methodology; Supervision; Writing - review & editing.

Edson Marchiori: Conceptualization; Data curation; Formal analysis; Investigation; Methodology; Supervision; Writing - review & editing.

Darcy Ribeiro Pinto Filho: Conceptualization; Data curation; Formal analysis; Investigation; Methodology; Supervision; Writing - review & editing.

Bruno Hochhegger: Conceptualization; Data curation; Formal analysis; Investigation; Methodology; Project administration; Resources; Supervision; Validation; Visualization; Writing - review & editing.

Declaration of competing interest

The authors declare that they have no conflict of interest.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.suronc.2021.101598>.

References

- [1] M. Luz, G. Strauss, D. Manzey, Impact of image-guided surgery on surgeons' performance: a literature review, *Int. J. Hum. Factors Ergon.* 4 (3–4) (2016) 229–263.
- [2] D. Manzey, S. Röttger, J.E. Bahner-Heyne, D. Schulze-Kissing, A. Dietz, J. Meixensberger, G. Strauss, Image-guided navigation: the surgeon's perspective on performance consequences and human factors issues, *Int. J. Med. Robot. Comput. Assist. Surg.* 5 (3) (2009) 297–308.
- [3] D.M. Dalgorf, R. Sacks, P.J. Wormald, Y. Naidoo, B. Panizza, B. Uren, C. Brown, J. Curotta, K. Snidvongs, R.J. Harvey, Image-guided surgery influences perioperative morbidity from endoscopic sinus surgery: a systematic review and meta-analysis, *Otolaryngology-Head Neck Surg.* (Tokyo) 149 (1) (2013) 17–29.
- [4] R.R. Casiano, W.A. Numa Jr., Efficacy of computed tomographic image-guided endoscopic sinus surgery in residency training programs, *Laryngoscope* 110 (8) (2000) 1277–1282.
- [5] M.C. Ingram, N. Atuegwu, L. Mawn, R.L. Galloway, Transorbital therapy delivery: phantom testing. In *Medical Imaging, Visualization, Image-Guided Procedures, and Modeling* 2011 Mar 1, vol. 7964, 2011, p. 79642A (International Society for Optics and Photonics).
- [6] S.K. Wise, R.J. Harvey, J.C. Goddard, P.O. Sheahan, R.J. Schlosser, Combined image guidance and intraoperative computed tomography in facilitating endoscopic orientation within and around the paranasal sinuses, *Am. J. Rhinol.* 22 (6) (2008) 635–641.
- [7] K. Stelter, B. Ertl-Wagner, M. Luz, S. Müller, G. Ledderose, V. Siedek, A. Berghaus, S. Arpe, A. Leunig, Evaluation of an image-guided navigation system in the training of functional endoscopic sinus surgeons. a prospective, randomised clinical study, *Rhinology* 49 (4) (2011) 429.
- [8] L. Xue, H. Fan, W. Shi, D. Ge, Y. Zhang, Q. Wang, Y. Yuan, Preoperative 3-dimensional computed tomography lung simulation before video-assisted thoroscopic anatomic segmentectomy for ground glass opacity in lung, *J. Thorac. Dis.* 10 (12) (2018) 6598.
- [9] K. Nakayama, Y. Oshiro, R. Miyamoto, K. Kohno, K. Fukunaga, N. Ohkohchi, The effect of three-dimensional preoperative simulation on liver surgery, *World J. Surg.* 41 (7) (2017) 1840–1847.
- [10] T. Kamai, N. Furuya, T. Kambara, H. Abe, M. Honda, Y. Shioyama, Y. Kaji, K. I. Yoshida, Single minimum incision endoscopic radical nephrectomy for renal tumors with preoperative virtual navigation using 3D-CT volume-rendering, *BMC Urol.* 10 (1) (2010 Dec 1) 7.
- [11] D. Wang, B. Zhang, X. Yuan, X. Zhang, C. Liu, Preoperative planning and real-time assisted navigation by three-dimensional individual digital model in partial nephrectomy with three-dimensional laparoscopic system, *International journal of computer assisted radiology and surgery* 10 (9) (2015 Sep 1) 1461–1468.
- [12] N. Ikeda, A. Yoshimura, M. Hagiwara, S. Akata, H. Saji, Three dimensional computed tomography lung modeling is useful in simulation and navigation of lung cancer surgery, *Ann. Thorac. Cardiovasc. Surg.* 19 (1) (2013 Feb 20) 1–5.
- [13] T. Akiba, Utility of three-dimensional computed tomography in general thoracic surgery, *General thoracic and cardiovascular surgery* 61 (12) (2013) 676–684.
- [14] K. Shimizu, S. Nakazawa, T. Nagashima, H. Kuwano, A. Mogi, 3D-CT anatomy for VATS segmentectomy, *J. Vis. Surg.* 3 (2017).
- [15] J. Hallet, B. Gayet, A. Tsung, G. Wakabayashi, P. Pessaux, 2nd International Consensus Conference on Laparoscopic Liver Resection Group, Systematic review of the use of pre-operative simulation and navigation for hepatectomy: current status and future perspectives, *J. Hepato-Biliary-Pancreatic Sci.* 22 (5) (2015) 353–362.
- [16] G. Quero, A. Lapergola, L. Soler, M. Shabaz, A. Hostettler, T. Collins, J. Marescaux, D. Mutter, M. Diana, P. Pessaux, Virtual and augmented reality in oncologic liver surgery, *Surgical Oncology Clinics* 28 (1) (2019 Jan 1) 31–44.
- [17] A. Zygomalas, D. Karavias, D. Koutsouris, I. Maroulis, D.D. Karavias, K. Giokas, V. Megalooikonomou, Computer-assisted liver tumor surgery using a novel semiautomatic and a hybrid semiautomatic segmentation algorithm, *Med. Biol. Eng. Comput.* 54 (5) (2016 May 1) 711–721.
- [18] F.J. Detmer, J. Hettig, D. Schindele, M. Schostak, C. Hansen, Virtual and augmented reality systems for renal interventions: a systematic review, *IEEE reviews in biomedical engineering* 10 (2017 Sep 6) 78–94.
- [19] F. Porpiglia, D. Amparore, E. Checcucci, R. Autorino, M. Manfredi, G. Iannizzi, C. Fiori, ESUT Research Group, Current use of three-dimensional model technology in urology: a road map for personalised surgical planning, *European urology focus* 4 (5) (2018 Sep 1) 652–656.
- [20] Z. Sun, D. Liu, A systematic review of clinical value of three-dimensional printing in renal disease, *Quant. Imag. Med. Surg.* 8 (3) (2018) 311.
- [21] A. Navaratnam, H. Abdul-Muhsin, M. Humphreys, Updates in urologic robot assisted surgery, *F1000Research* 7 (2018).
- [22] K. Slim, E. Nini, D. Forestier, F. Kwiatkowski, Y. Panis, J. Chipponi, Methodological index for non-randomized studies (MINORS): development and validation of a new instrument, *ANZ J. Surg.* 73 (9) (2003 Sep) 712–716.
- [23] G.M. Sullivan, R. Feinn, Using effect size—or why the P value is not enough, *Journal of graduate medical education* 4 (3) (2012) 279–282.
- [24] X. Wan, W. Wang, J. Liu, T. Tong, Estimating the sample mean and standard deviation from the sample size, median, range and/or interquartile range, *BMC Med. Res. Methodol.* 14 (1) (2014 Dec 1) 135.
- [25] J.P. Higgins, S.G. Thompson, Quantifying heterogeneity in a meta-analysis, *Stat. Med.* 21 (11) (2002) 1539–1558.
- [26] M. Egger, G.D. Smith, M. Schneider, C. Minder, Bias in meta-analysis detected by a simple, graphical test, *Br. Med. J.* 315 (7109) (1997 Sep 13) 629–634.
- [27] C.B. Begg, M. Mazumdar, Operating characteristics of a rank correlation test for publication bias, *Biometrics* (1994) 1088–1101.
- [28] T. Akiba, H. Marushima, J. Harada, S. Kobayashi, T. Morikawa, Importance of preoperative imaging with 64-row three-dimensional multidetector computed tomography for safer video-assisted thoracic surgery in lung cancer, *Surg. Today* 39 (10) (2009 Oct 1) 844–847.
- [29] X.B. Wei, J. Xu, N. Li, Y. Yu, J. Shi, W.X. Guo, H.Y. Cheng, M.C. Wu, W.Y. Lau, S. Q. Cheng, The role of three-dimensional imaging in optimizing diagnosis, classification and surgical treatment of hepatocellular carcinoma with portal vein tumor thrombus, *HPB* 18 (3) (2016 Mar 1) 287–295.
- [30] C.H. Fang, H.S. Tao, J. Yang, Z.S. Fang, W. Cai, J. Liu, Y.F. Fan, Impact of three-dimensional reconstruction technique in the operation planning of centrally located hepatocellular carcinoma, *J. Am. Coll. Surg.* 220 (1) (2015 Jan 1) 28–37.
- [31] Z. Wang, L. Qi, P. Yuan, X. Zu, W. Chen, Z. Cao, Y. Li, L. Wang, Application of three-dimensional visualization technology in laparoscopic partial nephrectomy of renal tumor: a comparative study, *J. Laparoendosc. Adv. Surg. Tech.* 27 (5) (2017 May 1) 516–523.
- [32] F. Porpiglia, C. Fiori, E. Checcucci, D. Amparore, R. Bertolo, Hyperaccuracy three-dimensional reconstruction is able to maximize the efficacy of selective clamping during robot-assisted partial nephrectomy for complex renal masses, *Eur. Urol.* 74 (5) (2018) 651–660.
- [33] J.D. Shirik, D.D. Thiel, E.M. Wallen, J.M. Linehan, W.M. White, K.K. Badani, J. R. Porter, Effect of 3-dimensional virtual reality models for surgical planning of robotic-assisted partial nephrectomy on surgical outcomes: a randomized clinical trial, *JAMA network open* 2 (9) (2019 Sep 4), e1911598.
- [34] D. Dindo, N. Demartines, P.-A. Clavien, Classification of surgical complications: a new proposal with evaluation in a cohort of 6336 patients and results of a survey, *Ann. Surg.* 240 (2) (2004) 205.
- [35] H. Cheng, B.P. Chen, I.M. Soleas, N.C. Ferko, C.G. Cameron, P. Hinoul, Prolonged operative duration increases risk of surgical site infections: a systematic review, *Surg. Infect.* 18 (6) (2017 Aug 1) 722–735.
- [36] E. Korol, K. Johnston, N. Waser, F. Sifakis, H.S. Jafri, M. Lo, M.H. Kyaw, A systematic review of risk factors associated with surgical site infections among surgical patients, *PLoS One* 8 (12) (2013 Dec 18), e83743.
- [37] C. Gibbons, C. Gibbons, Identification of Risk Factors by Systematic Review and Development of Risk-Adjusted Models for Surgical Site Infection, Prepress Projects Limited, 2011.
- [38] M.N. Mavros, S. Athanasiou, V.G. Alexiou, P.K. Mitsikostas, G. Peppas, M. E. Falagas, Risk factors for mesh-related infections after hernia repair surgery: a meta-analysis of cohort studies, *World J. Surg.* 35 (11) (2011 Nov 1) 2389.
- [39] N. Cornellà, J. Sancho, A. Sitges-Serra, Short and long-term outcomes after surgical procedures lasting for more than six hours, *Sci. Rep.* 7 (1) (2017) 1–8.
- [40] T. Catanzarite, S. Saha, M.A. Pilecki, J.Y. Kim, M.P. Milad, Longer operative time during benign laparoscopic and robotic hysterectomy is associated with increased 30-day perioperative complications, *J. Minim. Invasive Gynecol.* 22 (6) (2015 Sep 1) 1049–1058.
- [41] S. Singh, K. Swarer, K. Resnick, Longer operative time is associated with increased post-operative complications in patients undergoing minimally-invasive surgery for endometrial cancer, *Gynecol. Oncol.* 147 (3) (2017) 554–557.
- [42] T.D. Jackson, J.J. Wannares, R.T. Lancaster, D.W. Rattner, M.M. Hutter, Does speed matter? The impact of operative time on outcome in laparoscopic surgery, *Surg. Endosc.* 25 (7) (2011 Jul) 2288–2295, <https://doi.org/10.1007/s00464-010-1550-8>. Epub 2011 Feb 7. PMID: 21298533; PMCID: PMC3676304.
- [43] T. Fardoun, D. Chaste, E. Oger, R. Mathieu, B. Peyronnet, N. Rioux-Leclercq, G. Verhoest, J.J. Patard, K. Bensalah, Predictive factors of hemorrhagic complications after partial nephrectomy, *Eur. J. Surg. Oncol.* 40 (1) (2014 Jan 1) 85–89.
- [44] R. Gupta, D. Fuks, C. Bourdeaux, P. Radkani, T. Nomi, C. Lamer, B. Gayet, Impact of intraoperative blood loss on the short-term outcomes of laparoscopic liver resection, *Surg. Endosc.* 31 (11) (2017 Nov 1) 4451–4457.
- [45] X.P. Zhang, Y.Z. Gao, Z.H. Chen, K. Wang, Y.Q. Cheng, W.X. Guo, J. Shi, C. Q. Zhong, F. Zhang, S.Q. Cheng, In-hospital mortality after surgical resection in hepatocellular carcinoma patients with portal vein tumor thrombus, *J. Canc.* 10 (1) (2019) 72.
- [46] T. Helling, B. Blondeau, B. Wittek, Perioperative factors and outcome associated with massive blood loss during major liver resections, *HPB* 6 (3) (2004) 181–185.
- [47] S.C. Katz, J. Shia, K.H. Liao, M. Gonen, L. Ruo, W.R. Jarnagin, Y. Fong, M. I. D'Angelica, L.H. Blumgart, R.P. DeMatteo, Operative blood loss independently predicts recurrence and survival after resection of hepatocellular carcinoma, *Ann. Surg.* 249 (4) (2009 Apr 1) 617–623.
- [48] S. Li, K. Zhou, Y. Lai, C. Shen, Y. Wu, G. Che, Estimated intraoperative blood loss correlates with postoperative cardiopulmonary complications and length of stay in patients undergoing video-assisted thoracoscopic lung cancer lobectomy: a retrospective cohort study, *BMC Surg.* 18 (1) (2018 Dec 1) 29.

- [49] M. Rahouma, M. Kamel, G. Ghaly, A. Nasar, S. Harrison, B. Stiles, N. Altorki, J. L. Port, PS01. 45: intraoperative blood loss is an independent predictor of poor disease free survival for patients undergoing VATS lobectomy for lung cancer: topic: surgery, *J. Thorac. Oncol.* 11 (11) (2016 Nov 1) S297.
- [50] H. Nakamura, H. Saji, N. Kurimoto, T. Shinmyo, R. Tagaya, Impact of intraoperative blood loss on long-term survival after lung cancer resection, *Ann. Thorac. Cardiovasc. Surg.* 21 (1) (2015) 18–23, <https://doi.org/10.5761/atcs.0a.13-00312>. Epub 2014 Feb 28.
- [51] B.J. Shin, A.R. James, I.U. Njoku, R. Härtl, Pedicle screw navigation: a systematic review and meta-analysis of perforation risk for computer-navigated versus freehand insertion: a review, *J. Neurosurg. Spine* 17 (2) (2012 Aug 1) 113–122.
- [52] R. Bailer, R.C. Martin II, The effectiveness of using 3D reconstruction software for surgery to augment surgical education, *Am. J. Surg.* 218 (5) (2019 Nov 1) 1016–1021.
- [53] C.T. Yeo, A. MacDonald, T. Ungi, A. Lasso, D. Jalink, B. Zevin, G. Fichtinger, S. Nanji, Utility of 3D reconstruction of 2D liver computed tomography/magnetic resonance images as a surgical planning tool for residents in liver resection surgery, *J. Surg. Educ.* 75 (3) (2018 May 1) 792–797.