

Retrospective Cohort Study

Phase angle and non-alcoholic fatty liver disease before and after bariatric surgery

Joise Teixeira, Cláudio Augusto Marroni, Paula Rosales Zubiaurre, Ana Henz, Lais Faina, Lilian Kethelyn Pinheiro, Claudio Cora Mottin, Sabrina Alves Fernandes

ORCID number: Joise Teixeira 0000-0002-9267-5309; Cláudio Augusto Marroni 0000-0002-1718-6548; Paula Rosales Zubiaurre 0000-0002-9222-492X; Ana Henz 0000-0002-4260-2881; Lais Faina 0000-0003-2048-448X; Lilian Kethelyn Pinheiro 0000-0003-0402-634X; Claudio Cora Mottin 0000-0002-1638-2248; Sabrina Alves Fernandes 0000-0001-8504-603X.

Author contributions: Teixeira J and Fernandes SA performed the data collection; all the authors wrote the paper and approved the final version.

Institutional review board statement: This project was approved by the Research Ethics Committee of the Pontifícia Universidade Católica do Rio Grande do Sul (PUCRS). No. 2.423.466.

Informed consent statement: Patients who accepted to participate in the study signed the Informed Consent Form.

Conflict-of-interest statement: The authors declare no conflicts of interest for this article.

STROBE statement: The authors have read the STROBE Statement checklist of items, and the

Joise Teixeira, Department of Postgraduate Program in Medicine: Hepatology, Universidade Federal de Ciências da Saúde de Porto Alegre, Porto Alegre 90050170, RS, Brazil

Cláudio Augusto Marroni, Department of Gastroenterology and Hepatology, Universidade Federal de Ciências da Saúde de Porto Alegre (UFCSPA), Porto Alegre 91760470, RS, Brazil

Paula Rosales Zubiaurre, Department of Center of Morbid Obesity, Pontifícia Universidade Católica do Rio Grande do Sul (PUCRS), Porto Alegre 90610000, RS, Brazil

Ana Henz, Lilian Kethelyn Pinheiro, Sabrina Alves Fernandes, Department of Nutrition, Centro Universitário Metodista (IPA), Porto Alegre 90420060, RS, Brazil

Lais Faina, Department of Vascular Surgery, Hospital Federal dos Servidores do Estado do Rio de Janeiro, Rio de Janeiro 20221161, RJ, Brazil

Claudio Cora Mottin, Department of Obesity and Metabolic Syndrome Center, Pontifícia Universidade Católica do Rio Grande do Sul (PUCRS), Porto Alegre 90619-900, RS, Brazil

Corresponding author: Sabrina Alves Fernandes, RN, Research Scientist, Department of Nutrition, Centro Universitário Metodista (IPA), Coronel Joaquim Pedro Salgado, Porto Alegre 90420060, RS, Brazil. sabrina@drasabrinafernandes.com

Abstract

BACKGROUND

Obesity is a global health problem that is continuing to increase in the young population. In Brazil, the frequency of obesity in 2018 was 19.8%. Several comorbidities are directly associated with obesity, such as non-alcoholic fatty liver disease (NAFLD), which is considered the most common liver disorder in Western countries and affects up to 46% of adults. Bariatric surgery is effective in treating obesity and can improve NAFLD; however, the effect of bariatric surgery on body composition, phase angle (PA), and improving NAFLD needs to be further studied.

AIM

To analyze the PA in the postoperative period of bariatric surgery and to correlate it with changes in body composition and liver disease.

manuscript was prepared and revised according to the STROBE Statement checklist of items.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Invited manuscript

Specialty type: Gastroenterology and hepatology

Country/Territory of origin: Brazil

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): 0
Grade C (Good): C
Grade D (Fair): 0
Grade E (Poor): 0

Received: June 30, 2020

Peer-review started: June 30, 2020

First decision: August 8, 2020

Revised: August 24, 2020

Accepted: October 12, 2020

Article in press: October 12, 2020

Published online: November 27, 2020

P-Reviewer: Sasaki A

S-Editor: Fan JR

L-Editor: Filipodia

P-Editor: Xing YX



METHODS

This study is a retrospective cohort study of the analysis of the medical records of patients undergoing bariatric surgery in a reference center of a teaching hospital in Porto Alegre over a 2-year period. Patients older than 18 years whose record contained all information relevant to the study were included. The data analyzed were body composition and PA through electrical bioimpedance and NAFLD through liver biopsy in the pre- and postoperative period. The level of significance adopted for the statistical analyses was 5%.

RESULTS

We evaluated 379 patients with preoperative data. Regarding PA, 169 patients were analyzed, and 33 patients had liver biopsy pre- and postoperatively with NAFLD information. In total, 79.4% were female, with a mean age of 39.1 ± 10.6 years. The average body mass index (BMI) was 45.9 ± 7.5 kg/m². The PA showed a mean of $5.8 \pm 0.62^\circ$ in the preoperative period and a significant reduction in the postoperative period. A postoperative reduction in body composition data (skeletal muscle mass, fat percentage, fat mass, body cell mass, BMI and visceral fat area) was shown as well. Regarding liver disease, all patients presented a reduction in the degrees and stages of liver disease in the postoperative period, and some had no degree of liver disease at all.

CONCLUSION

PA decreased after bariatric surgery, with a direct correlation with weight loss and changes in body composition. The decrease in PA was not correlated with the improvement in NAFLD.

Key Words: Obesity; Body composition; Bariatric surgery; Phase angle; Non-alcoholic fatty liver disease; Liver disease

©The Author(s) 2020. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: We retrospectively evaluated 379 patients who underwent bariatric surgery, with non-alcoholic fatty liver disease in the preoperative period; we compared body composition, phase angle (PA) behavior and change in non-alcoholic fatty liver disease (NAFLD) in the pre- and postoperative period. There was an important improvement in body composition/body fat percentage and an improvement in NAFLD after bariatric surgery. Worsening PA was directly correlated with weight loss and skeletal muscle mass.

Citation: Teixeira J, Marroni CA, Zubiaurre PR, Henz A, Faina L, Pinheiro LK, Mottin CC, Fernandes SA. Phase angle and non-alcoholic fatty liver disease before and after bariatric surgery. *World J Hepatol* 2020; 12(11): 1004-1019

URL: <https://www.wjgnet.com/1948-5182/full/v12/i11/1004.htm>

DOI: <https://dx.doi.org/10.4254/wjh.v12.i11.1004>

INTRODUCTION

Obesity is a complex chronic inflammatory disease characterized by excessive accumulation of body fat. It has multiple etiologies, such as biochemical, genetic, behavioral, social, environmental, and nutritional factors and imbalance between food intake and energy expenditure^[1-4]. This disease presents its own pathophysiology as well as associated comorbidities^[5].

According to VIGITEL Brazil 2018^[6], the frequency of obese adults was 19.8% in a society that is aging and more ill. Obesity is closely associated with cardiovascular diseases, dyslipidemia, fatty liver, diabetes mellitus (DM) and other endocrine and metabolic disorders^[1]. In obese people, a common comorbidity is non-alcoholic fatty liver disease (NAFLD).

NAFLD is defined as accumulation of fat in more than 5% of hepatocytes^[7] without secondary cause, such as alcohol consumption (> 20 g for women and > 30 g for men,

daily), use of steatogenic medications, or hereditary disorders^[8].

The gold standard for the diagnosis of NAFLD is liver biopsy^[9], and its classification encompasses a wide spectrum of histopathological changes, from simple hepatic steatosis, which can evolve from non-alcoholic steatohepatitis (NASH), to cirrhosis and/or hepatocellular carcinoma^[10]. Therefore, NAFLD, due to its different staging, may or may not present fibrosis in hepatocytes^[11,12]. NAFLD is considered the most common liver disease in Western countries, affecting 90% of morbidly obese patients eligible for bariatric surgery^[11].

Because of the difficulty found in the clinical treatment of obesity, bariatric surgery is more efficient as a treatment option for individuals with severe obesity, when compared to non-surgical interventions^[13]. Furthermore, surgical treatment shows improvement or remission in NAFLD^[14].

According to the Brazilian Society of Bariatric and Metabolic Surgery^[15], bariatric surgery combines techniques aimed to treat morbid obesity, severe obesity, and diseases associated with excess body fat or exacerbated by it. Gastric bypass surgery in Y by Roux (BPGYR) or Fobi-Capella surgery is the most commonly performed technique in Brazil and worldwide.

Electrical bioimpedance (BIA) is a method for analyzing body composition and is based on the principle of resistance and reactance that cells impose on the electrical current emitted by the device^[16]. The human body is constituted of conductors like water and non-conductors, like body fat^[17,18].

Several parameters are measured using BIA, including body water, lean mass, fat mass, and phase angle (PA). BIA assesses nutritional status and can be a good method for prognostic evaluation, as it is practical, fast, non-invasive, and low cost^[19]. However, body composition values in patients with dysmorphia (edema, ascites, and morbid obesity), as measured by BIA, may suffer interference. This is why PA has been widely used, since it is not associated with interference^[20,21].

Currently, there are segmented, multifrequency BIA devices with greater precision for assessing body composition in morbidly obese patients, as validated by Faria *et al*^[22].

PA was originally described by Baumgartner *et al*^[23] for the diagnosis of metabolic disorders. It is a parameter applicable in clinical practice because it reliably helps to describe cell vitality and integrality. High values (up to 8°) may indicate body homeostasis, whereas values below 6°, depending on the disease, reflect a poor clinical prognosis, indicating changes in the selective permeability of the cell membrane^[23,24].

As already mentioned, PA reflects cellular integrity and functionality by measuring, through an electrical current, the values of resistance and reactance of the membrane of these cells, with skeletal muscle being a conductor of electrical current and the opposite occurring with fat mass^[23-25]. In view of this fact, we believe that the body change resulting from bariatric surgery will reflect an improvement in NAFLD and can be measured by PA.

To date, there are not enough studies evaluating morbid obesity, body composition (described by the BIA), and associated comorbidities, such as NAFLD.

The present study aims to analyze the behavior of PA in the postoperative period of bariatric surgery, correlating it with changes in body composition and improvement of liver disease.

MATERIALS AND METHODS

This is a retrospective cohort study that analyzed the medical records of patients undergoing bariatric surgery in a referral center of a teaching hospital in Porto Alegre. Patients over 18-years-old whose record contained all the information relevant to the study in the electronic or physical medical record were included. Patients who did not contain complete data were excluded. For convenience, the sample was carried out from July 2015 to July 2017. The data obtained were related to the protocol for routine pre- and postoperative care at the service's outpatient clinic.

Electrical bioimpedance-PA and body composition

Data on body composition and weight were measured using BIA in all patients in the week preceding bariatric surgery without prior preparation. For BIA, the patient stood upright on the InBody 770 device from Ottoboni, with an electric current intensity of 80 μ A and 50/60 kHz frequency. The PA was obtained through the values of resistance and reactance through the formula: $PA = \text{tangent arc } (Xc/R) \times 180/3.1416$, described in the result sheet. The BIA was performed in a second step, and for

comparative postoperative analysis, we analyzed those patients who used the same BIA device 6 mo to 12 mo after bariatric surgery for routine postoperative evaluation.

Anthropometric data-height and body mass index

Height was measured using a wall Tonelli stadiometer, model E150 A, with the patient standing upright and barefoot, with their feet together, and with their backs positioned against the wall. Body mass index (BMI) was calculated using the equation weight in kilograms, divided by height in meters squared and classified according to the World Health Organization^[4]: BMI ≥ 30 kg/m² to 34.9 kg/m² Obesity Grade I, BMI between 35 kg/m² to 39.9 kg/m² Obesity Grade II, and BMI ≥ 40 kg/m² Obesity Grade III. The group of patients with a BMI greater than 50 kg/m² was analyzed separately.

Bariatric surgery

The bariatric surgery used was the BPGYR with intestinal derivation after monitoring and preparation with a multidisciplinary service team. All patients met the criteria for bariatric surgery. The surgeries were performed by four specialist surgeons, from the same team, trained and with much experience.

Liver biopsy-NAFLD

Liver biopsies were routinely performed during the bariatric surgery trans operation by the surgeon under direct vision using a Tru-Cut needle at the beginning of the surgical procedure before liver withdrawal. The biopsies were analyzed by the same pathologist at the Hospital's Pathology Laboratory. The classification was made using the criteria of Kleiner *et al*^[26], as follows: Absence or presence of NAFLD and/or cirrhosis; steatosis activity (absent, discreet, moderate, accented, and massive-according to grade and location of the injury); ballooning and lobular inflammation; degrees of NASH (1, 2, 3, and 4), and fibrosis stages (1, 2, 3 and 4). The liver biopsies performed in the postoperative period were obtained by the same surgical team in patients who underwent a second intervention (cholecystectomy or appendectomy) in the period from 6 to 12 mo after bariatric surgery and analyzed in the same way as those of the first biopsy.

Statistical analysis

Quantitative variables are described as mean and standard deviation or median and interquartile range. Categorical variables are described by absolute and relative frequencies.

To compare means before and after bariatric surgery, the t-student test for paired samples was applied. When comparing nominal categorical variables, the McNemar test was used and, for ordinals, the Wilcoxon test. To compare means between genders, the t-student test for independent samples was applied.

In the association between quantitative and ordinal variables, Pearson or Spearman correlation tests were used.

The level of significance adopted was 5% ($P < 0.05$), and the analyses were performed using the SPSS version 21.0 program (Armonk, NY, United States).

The project was elaborated in accordance with resolution 466 of 2012, which regulates the conduct of research in human beings, and submitted to and approved by the Research Ethics Committee under number 2.423.466. Patients who accepted to participate in the study signed the Informed Consent Term.

This study was reviewed by our specialist Biostatistics, Mestre, Ceres Andréia Vieira de Oliveira.

RESULTS

Of the 727 patients operated on in the period, 379 who had complete preoperative information were allocated. For the analysis of data related to PA, 169 of these patients who underwent pre- and post-evaluation on the same device were allocated. Regarding NAFLD, we analyzed 33 patients who underwent postoperative liver biopsy.

Of the 379 patients, 79.4% were female, with a mean age of 39.1 ± 10.6 years and a BMI of 45.9 ± 7.5 kg/m², classified as Obesity Grade III^[4]. It is noteworthy that more than 22% of patients had a BMI greater than 50 kg/m². Full-body PA showed an average of $5.89^\circ \pm 0.62^\circ$, with a minimum of 4.3° and a maximum of 7.9° . The other characteristics of the sample are shown in [Table 1](#).

Table 1 Sample characteristics, *n* = 379

Variables	<i>n</i> = 379
Gender, <i>n</i> (%)	
Male	78 (20.6)
Female	301 (79.4)
Age in yr, mean ± SD	39.1 ± 10.6
Weight in kg, mean ± SD	123.7 ± 24.8
Minimum	77
Maximum	235
BMI in kg/m ² , mean ± SD	45.9 ± 7.5
BMI classification, <i>n</i> (%)	
30-34.99 kg/m ²	3 (0.8)
35-39.99 kg/m ²	82 (21.6)
40-49.99 kg/m ²	210 (55.4)
50 kg/m ² or more	84 (22.2)
SMM in kg, mean ± SD	34.1 ± 7.5
BCM in kg, mean ± SD	39.4 ± 8.3
Fat mass in kg, mean ± SD	62.6 ± 15.1
% Fat-mean ± SD	50.7 ± 4.6
Visceral fat area in cm ² , mean ± SD	243.7 ± 31.2
Phase angle °, mean ± SD	
Full-body	5.89 ± 0.62
RA	5.58 ± 0.62
LA	5.42 ± 0.65
Tr	7.97 ± 1.18
RL	6.14 ± 0.81
LL	6.07 ± 0.84

BCM: Body cell mass; BMI: Body mass index; LA: Left arm; LL: Left leg; RA: Right arm; RL: Right leg; SD: Standard deviation; SMM: Skeletal muscle mass; Tr: Torso.

All patients (*n* = 379) were diagnosed with NAFLD by liver biopsy, and the histological characteristics are shown in Table 2. Regarding the degrees of the disease, 78.1% had NASH and 43% fibrosis, ranging from F1 to F3. No patient had cirrhosis.

The difference in pre- and postoperative body composition (*n* = 379) is found in Table 3, with all items statistically significant (*P* < 0.001).

The associations of body composition variables with PA, stratified by gender (*n* = 379), are shown in Table 4. In males, there was an inverse, statistically significant association between the percentage of fat, fat mass, and weight with PA in most body compartments, except the torso.

In females, there was a negative, statistically significant association between the percentage of fat and PA in all compartments. There was a positive, statistically significant association between skeletal muscle mass (SMM) and body cell mass (BCM) with PA in most body compartments, except the legs.

There was a negative, statistically significant association (*r* = -0.483; *P* < 0.001) between the reduction of full-body PA and the percentage of weight loss, as shown in Figure 1. There was a statistically significant positive association between the reduction of full-body PA with loss of SMM (*r* = 0.307; *P* < 0.001), as shown in Figure 2. The association of the reduction in PA of the whole body was also positive and significant with the loss of fat mass, (*r* = 0.280; *P* < 0.001), MCC (*r* = 0.287; *P* < 0.001), visceral fat area (*r* = 0.275; *P* < 0.001), BMI (*r* = 0.413; *P* < 0.001), and variation in the fat

Table 2 Sample characteristics for liver disease, *n* = 379

Variables	<i>n</i> (%)
NAFLD	
Yes	379 (100.0)
No	0 (0.0)
Hepatic steatosis	
Absent	0 (0.0)
Discreet	159 (42.0)
Moderate	104 (27.4)
Accented	84 (22.2)
Massive	32 (8.4)
Ballooning	
Absent	91 (24.0)
Discreet	221 (58.3)
Moderate	18 (4.7)
Accented	49 (12.9)
Massive	0 (0.0)
Lobular inflammation	
Absent	243 (64.1)
Discreet	100 (26.4)
Moderate	28 (7.4)
Accented	8 (2.1)
Massive	0 (0.0)
NASH	
Absent	83 (21.9)
Grade 1	201 (53.0)
Grade 2	67 (17.7)
Grade 3	28 (7.4)
Grade 4	0 (0.0)
Absent	216 (57.0)
Grade 1	86 (22.7)
Grade 2	44 (11.6)
Grade 3	33 (8.7)
Grade 4	0 (0.0)
Cirrhosis	
Yes	0 (0.0)
No	379 (100.0)

NAFLD: Non-alcoholic fatty liver disease; NASH: Non-alcoholic steatohepatitis.

percentage ($r = 0.304$; $P < 0.001$), as shown in [Figure 3](#).

Men showed a more marked and significant reduction in the percentage of fat (-16 ± 4.9 vs -12.3 ± 5.3 ; $P < 0.001$) and visceral fat area (-110.7 ± 58.9 vs -90 ± 42.8 ; $P = 0.024$) when compared to women.

After bariatric surgery, a significant reduction in PA values ($n = 169$) was observed in all body compartments ($P < 0.001$), as described in [Table 5](#).

Table 3 Variation of body composition before and after bariatric surgery, *n* = 379

Variables	Pre	Post	P value
	mean ± SD	mean ± SD	
BMI in kg/m ²	45.9 ± 7.5	31.2 ± 5.0	< 0.001
SMM in kg	34.1 ± 7.5	29.0 ± 6.4	< 0.001
BCM in kg	39.4 ± 8.3	33.9 ± 6.9	< 0.001
Fat mass in kg	62.6 ± 15.1	31.9 ± 11.1	< 0.001
% Fat	50.7 ± 4.6	37.2 ± 8.1	< 0.001
Visceral fat area in cm ²	243.7 ± 31.2	152.0 ± 51.0	< 0.001

BCM: Body cell mass; BMI: Body mass index; SD: Standard deviation; SMM: Skeletal muscle mass.

Table 4 Association of body composition variables with phase angle in men and women, using Pearson's correlation coefficient, *n* = 379

Variables	Weight in kg	SMM in kg	BCM in kg	Fat mass in kg	% Fat	Visceral fat area in cm ²
Male gender, <i>n</i> = 78						
PA full body	-0.418 ^c	-0.118	-0.101	-0.469 ^c	-0.401 ^c	-0.075
PA RA	-0.299 ^b	-0.083	-0.064	-0.337 ^b	-0.287 ^a	-0.095
PA LA	-0.286 ^a	-0.031	-0.008	-0.349 ^b	-0.341 ^b	-0.136
PA Tr	-0.074	0.074	0.094	-0.134	-0.185	0.311 ^b
PA RL	-0.493 ^c	-0.179	-0.170	-0.531 ^c	-0.421 ^c	-0.101
PA LL	-0.509 ^c	-0.211	-0.209	-0.534 ^c	-0.403 ^c	-0.066
Female gender, <i>n</i> = 301						
PA full body	-0.106	0.125 ^a	0.122 ^b	-0.152 ^b	-0.393 ^c	-0.098
PA RA	0.011	0.169 ^b	0.148 ^a	-0.036	-0.245 ^c	-0.111
PA LA	0.043	0.193 ^b	0.155 ^b	-0.009	-0.226 ^c	-0.135 ^b
PA Tr	0.024	0.120 ^a	0.133 ^a	-0.030	-0.172 ^b	-0.058
PA RL	-0.154 ^b	0.083	0.068	-0.182 ^b	-0.406 ^c	-0.043
PA LL	-0.151 ^b	0.069	0.052	-0.187 ^b	-0.388 ^c	-0.086

^a*P* < 0.05.

^b*P* < 0.01.

^c*P* < 0.001. BCM: Body cell mass; LA: Left arm; LL: Left leg; PA: Phase angle; RA: Right arm; RL: Right leg; SMM: Skeletal muscle mass; Tr: Torso.

There was a statistically significant inverse association between BMI and full-body PA (*P* < 0.018), right leg and left leg (*P* < 0.001), as shown in Table 6. Regarding PA and NAFLD, there was no significant association (*P* > 0.05).

The analysis of the postoperative liver biopsy, compared with the preoperative biopsy of these patients (*n* = 33), showed that all of them had a reduction in the degrees and stages of liver disease, and 18.2% had no degree of liver disease (*P* < 0.05).

The other histological changes, which decreased from 75 to 90%, are described in Table 7. The body composition of this group (*n* = 33) showed that all parameters significantly decreased (*P* < 0.001) and that there was a reduction in PA in all compartments (*P* < 0.001), as described in Table 8.

There was a positive, statistically significant association between BMI (kg/m²) and lobular inflammation (*P* < 0.05), NASH (*P* < 0.01), and fibrosis (*P* < 0.05). The other body composition variables did not correlate with the different histological characteristics of NAFLD.

There was an association between the variations of PA before and after bariatric surgery regarding the degree of lobular inflammation (*r*_s = -0.593; *P* = < 0.001) but not

Table 5 Variation of phase angle in the pre- and postoperative period of bariatric surgery, *n* = 169

Variables	Pre	Post	P value
	mean ± SD	mean ± SD	
PA full body	5.92 ± 0.55	4.98 ± 0.55	< 0.001
PA RA	5.58 ± 0.60	4.73 ± 0.59	< 0.001
PA LA	5.42 ± 0.63	4.54 ± 0.57	< 0.001
PA Tr	7.87 ± 1.15	6.76 ± 1.29	< 0.001
PA RL	6.25 ± 0.66	5.19 ± 0.65	< 0.001
PA LL	6.18 ± 0.70	5.14 ± 0.66	< 0.001

LA: Left arm; LL: Left leg; PA: Phase angle; RA: Right arm; RL: Right leg; SD: Standard deviation; Tr: Torso.

Table 6 Association of body mass index with phase angle through Pearson correlation, *n* = 169

Variables	BMI	P value
	<i>r</i>	
PA full body	-0.121	0.018
PA RA	0.031	0.551
PA LA	0.057	0.265
PA Tr	0.054	0.298
PA RL	-0.262	< 0.001
PA LL	-0.262	< 0.001

BMI: Body mass index; LA: Left arm; LL: Left leg; PA: Phase angle; *r*: Pearson's correlation coefficient; RA: Right arm; RL: Right leg; Tr: Torso.

steatosis ($r_s = 0.305$; $P = 0.095$), ballooning ($r_s = 0.057$; $P = 0.760$), NASH ($r_s = -0.197$; $P = 0.288$), and fibrosis ($r_s = -0.183$; $P = 0.324$).

DISCUSSION

Obese patients, candidates for bariatric surgery diagnosed with NAFLD, have been extensively studied^[8,27-31]; and, to date, changes in lifestyle are the only effective forms of treatment for NAFLD. It was established that a loss of 7%-10% of body weight is necessary to present any change^[32]. However, in order to guarantee body homeostasis, it is essential that the weight loss of these patients is the highest possible percentage of fat mass and not of SMM, preserving muscle volume and functionality. Therefore, BIA and PA are essential tools for monitoring the body characteristics and health of these patients^[19,33].

Most of our patients were women, with a mean age of 39.1 ± 10.6 years and a BMI of 45.9 ± 7.5 kg/m², findings similar to those of Losekann *et al*^[9], who in 2013 analyzed 250 patients with liver biopsies performed in the bariatric surgery trans operation, which 80% were women, with a mean age of 36.8 ± 10.2 years and a BMI of 43.6 ± 5.2 kg/m².

BMI is an analytical, non-laboratory method that is easy to apply and reproducible, allowing an indirect assessment of body composition, and is a defining parameter of indication for bariatric surgery.

The mean BMI after surgery decreased to 31.2 ± 5.0 kg/m² ($P < 0.001$), changing from Grade III Obesity to Grade I. The reduction in BMI was significant in both sexes, with no significant difference between them. All body composition parameters had a significant decrease ($P < 0.001$), mainly on fat mass and visceral fat area. In the present study, men compared to women showed a more marked and significant reduction in the percentage of fat and in the area of visceral fat.

Perrone *et al*^[34] reported that the decrease in BMI after bariatric surgery in men was

Table 7 Degrees and staging of non-alcoholic fatty liver disease before and after bariatric surgery, *n* = 33

Variables	Pre	Post	P value
	<i>n</i> (%)	<i>n</i> (%)	
NAFLD			0.031
Yes	33 (100)	27 (81.8)	
No	0 (0.0)	6 (18.2)	
Hepatic steatosis			< 0.001
Absent	0 (0.0)	6 (18.2)	
Discreet	14 (42.4)	24 (72.7)	
Moderate	8 (24.2)	3 (9.1)	
Accented	8 (24.2)	0 (0.0)	
Massive	3 (9.1)	0 (0.0)	
Ballooning			< 0.001
Absent	5 (15.2)	29 (87.9)	
Discreet	23 (69.7)	4 (12.1)	
Moderate	3 (9.1)	0 (0.0)	
Accented	2 (6.1)	0 (0.0)	
Lobular inflammation			0.003
Absent	21 (63.6)	30 (90.9)	
Discreet	7 (21.2)	3 (9.1)	
Moderate	3 (9.1)	0 (0.0)	
Accented	2 (6.1)	0 (0.0)	
NASH			< 0.001
Absent	6 (18.2)	25 (75.8)	
Grade 1	20 (60.6)	5 (15.2)	
Grade 2	4 (12.1)	3 (9.1)	
Grade 3	3 (9.1)	0 (0.0)	
Fibrosis			0.033
Absent	26 (78.8)	27 (81.8)	
Grade 1	3 (9.1)	4 (12.1)	
Grade 2	0 (0.0)	0 (0.0)	
Grade 3	4 (12.1)	2 (6.1)	

NAFLD: Non-alcoholic fatty liver disease; NASH: Non-alcoholic steatohepatitis.

greater than that in women, without influencing the improvement of comorbidities in the long term, because the BMI does not differentiate or qualify weight loss, which should be mostly fat and not SMM. A study by Hartwig *et al*^[35], showed similar results in men and women regarding the decrease in the percentage of fat and fat mass in the postoperative period. De Paris *et al*^[36] showed a reduction in body composition (weight, fat-free mass, SMM, fat mass, and fat percentage) in the postoperative period of bariatric surgery, results similar to ours.

All of our patients had NAFLD; accentuated or massive steatosis in 30.6%; ballooning in 76%; lobular inflammation in 35.9%; NASH in 78.1%; fibrosis in 43%; and no case of cirrhosis. NAFLD patients are obese^[37], with a prevalence of 51% of cases^[38]. In patients with obesity undergoing bariatric surgery, the percentage of steatosis ranged from 87.6% to 100%^[8,27-31], ballooning from 58.9% to 88%^[29,30], lobular inflammation from 23% to 88%^[29,31], and fibrosis from 31% to 44.9%^[28,29,31], findings

Table 8 Phase angle and body composition in patients who underwent liver biopsy after bariatric surgery, *n* = 33

Variables	Pre	Post	P value
	mean \pm SD	mean \pm SD	
BMI in kg/m ²	45.2 \pm 6.9	31.8 \pm 5.4	< 0.001
SMM in kg	31.9 \pm 6.8	27.1 \pm 5.3	< 0.001
BCM in kg	37.0 \pm 7.8	32.1 \pm 5.9	< 0.001
Fat mass in kg	60.7 \pm 13.3	33.6 \pm 11.1	< 0.001
% Fat	51.6 \pm 3.5	39.8 \pm 8.2	< 0.001
Visceral fat area in cm ²	249.3 \pm 23.1	162.5 \pm 52.3	< 0.001
PA full body °	5.93 \pm 0.65	4.91 \pm 0.60	< 0.001
PA RA °	5.55 \pm 0.77	4.61 \pm 0.64	< 0.001
PA LA °	5.44 \pm 0.86	4.44 \pm 0.58	< 0.001
PA Tr °	7.84 \pm 1.24	6.73 \pm 1.03	< 0.001
PA RL °	6.42 \pm 0.72	5.18 \pm 0.68	< 0.001
PA LL °	6.38 \pm 0.74	5.19 \pm 0.66	< 0.001

BCM: Body cell mass; BMI: Body mass index; LA: Left arm; LL: Left leg; PA: Phase angle; RA: Right arm; RL: Right leg; SMM: Skeletal muscle mass; Tr: Torso.




Figure 1 Association between the percentage of weight loss with the reduction of the phase angle in the postoperative period of bariatric surgery (*n* = 169).

similar to ours.

Some NAFLD patients present progression from simple steatosis to advanced stages, such as NASH and fibrosis, increasing the risk of cirrhosis and hepatocellular carcinoma. In addition, it is believed that NAFLD is implicated in the pathogenesis of type 2 DM and cardiovascular diseases^[32]. These facts are fundamental in the search for the reduction of obesity that bariatric surgery provides.

We found a direct association between BMI and lobular inflammation, NASH, and fibrosis (*P* < 0.05), which is why it is important to monitor evolution of this group of patients.

The assessment of body composition is limited in several clinical conditions; and, therefore, the use of BIA data has gained increasing attention^[39]. PA can be used as a biomarker for lean mass and/or for reducing muscle mass^[39-41], besides being recognized as a marker of malnutrition^[40,41] and predictor of morbidity and mortality in several diseases^[18,42].

The PA of healthy individuals can vary between 6° and 7°, according to Bosy-Westphal *et al*^[43] and 6.96° according to Barbosa-Silva *et al*^[44]. There is no reference value that classifies PA for patients with obesity who have NAFLD. In the studied population, the PA was 5.89 \pm 0.62°, lower than the values described above, probably




Figure 2 Association between the variation in skeletal muscle mass with the reduction of phase angle in the postoperative period of bariatric surgery ($n = 169$).




Figure 3 Association between the reduction in the percentage of body fat with the reduction in phase angle in the postoperative period of bariatric surgery ($n = 169$).

due to obesity.

In [Table 4](#), we evaluated the associations of PA with the variables of body composition, and we observed in men a significant negative relation with decreased weight, fat mass, and percentage of fat ($P < 0.01$). In women, this relation of PA was with the decrease in the percentage of fat ($P < 0.01$). There was a significant positive relation with SMM and BCM ($P < 0.05$). PA decreased in most variables related to weight loss ($P < 0.05$). Baumgartner *et al*^[45], in 1988, observed a statistically significant negative correlation of PA with the percentage of body fat, corroborating the findings of our study. A study by Peres *et al*^[46] analyzed 66 patients over 18 years of age with NAFLD, chronic hepatitis, cirrhosis, and hepatocellular carcinoma who had a mean PA of 5.1°, values similar to those found in the present study.

The post-surgical evaluation showed that PA decreased significantly in all evaluated segments ($P < 0.001$), as shown in [Table 5](#), and its correlation with BMI showed a significant difference ($P = 0.018$), as shown in [Table 6](#). These are apparently paradoxical findings, based on knowledge that the highest PA means improvement and the lowest means clinical worsening. We must consider the time of the second assessment, within the first year, as directly related to the decrease. A new evaluation of PA, at a longer time point, already programmed, may show an increase in PA.

Norman *et al*^[47] described that BMI is one of the biological factors that influences PA. Furthermore, Llamas *et al*^[33] found a reduction in PA in individuals with a BMI greater than 35 kg/m². Bosty-Westphal *et al*^[43] showed that PA increased with an increase in BMI up to 30 kg/m² and that such physiological behavior can be explained as a reflection of the increase in the number of cells (adipocytes and myocytes), since the reactivity of BIA is dependent on the amount of cell membranes. The same study

showed that with BMI above 40 kg/m², there was an inverse relation between PA and BMI. The explanation is that severe obese individuals have a loss of functionality in the cell membrane, which may contribute to the decrease in PA.

In 2017, Vassilev *et al*^[48] in Germany, with 173 patients undergoing bariatric surgery (BPGYR), showed a correlation between PA and weight loss, between 6 and 12 mo postoperatively, and a significant reduction in lean mass with a reduction in PA, results similar to ours. Koehler *et al*^[49], with 20 patients undergoing bariatric surgery (BPGYR), showed significant results in reducing PA, at an earlier time, from 3 to 6 mo.

In the present study, there was a negative association between the reduction of full-body PA and the percentage of weight loss ($P < 0.001$), as shown in **Figure 1**. There was a positive association between the reduction of full-body PA with the variation of SMM ($P < 0.001$), as shown in **Figure 2**; and there was a positive association between the reduction of PA of the whole body and the percentage variation of fat ($P < 0.001$), as shown in **Figure 3**. These same associations were observed in the other parameters (loss of fat mass; BCM; visceral fat area, BMI), all with significant value ($P < 0.001$).

In the Vassilev *et al* study^[48], the higher the percentage of body fat in the postoperative period, the lower the PA, and the reduction in BCM occurred with a reduction in PA after 9 mo of surgery. Thus, it is clear that the percentage of fat, even after weight loss, continues to have a negative influence on PA. The decrease in PA in the postoperative period is linked to weight loss, which can be a confusing factor when relating the reduction in the percentage of fat and PA, since the present study also associated a reduction in BMI with a decrease in PA.

There was an association between the variations of PA before and after bariatric surgery regarding the degree of lobular inflammation ($r_s = -0.593$; $P < 0.001$) but not with steatosis ($r_s = 0.305$; $P = 0.095$), ballooning ($r_s = 0.057$; $P = 0.760$), NASH ($r_s = -0.197$; $P = 0.288$), and fibrosis ($r_s = -0.183$; $P = 0.324$).

Regarding the role of PA and NAFLD staging, Peres *et al*^[46] found no difference in PA values in patients with different degrees and stages of liver disease; our findings did not show significance associating the decrease in PA with the variation in the degree of steatosis, ballooning, NASH, and fibrosis in the pre- and postoperative period, demonstrating that the improvement of liver disease after bariatric surgery is not related to the worsening of PA.

The comparison of liver histology before and after surgery ($n = 33$) showed a significant improvement in NAFLD, where 18.2% had no degree of liver disease ($P = 0.031$); all parameters analyzed showed a reduction in histological changes, from 75% to 90% ($P < 0.05$), as described in **Table 7**.

Máttar *et al*^[24] analyzed patients undergoing different bariatric surgery techniques and observed significant improvement in steatosis (from 88% to 8%), lobular inflammation (from 23% to 2%), and fibrosis (31% to 13%). In addition, 37% no longer had lobular inflammation, and 20% had no fibrosis, in line with our study. Similar data were found in Cazzo *et al*^[50] review in 2017 with patients undergoing bariatric surgery using different surgical techniques^[27,30,51-53].

A pioneering study by Silverman *et al*^[54] in 1995 observed an improvement in liver disease in the postoperative period of bariatric surgery with the same surgical technique we used, with a 71% reduction in steatosis; 19% absence of steatosis; 76.9% absence of fibrosis, and 7.6% reduction in fibrosis, corroborating with our findings. There was a marked change in body composition in this group ($n = 33$), in all parameters ($P < 0.001$), as well as a reduction in PA ($P < 0.001$), as shown in **Table 8**. These data reinforce that weight loss is fundamental in the treatment of NAFLD.

CONCLUSION

The PA decreased after bariatric surgery, with a direct correlation with weight loss and changes in body composition. Grade III obesity became Grade I obesity. The decrease in the PA after bariatric surgery did not correlate with the improvement of NAFLD, even with the marked improvement of NAFLD. The decrease in the PA after performing bariatric surgery correlated with the decrease in the BMI, loss of SMM, and decreases in body fat (in percentages and kilograms), BCM, and visceral fat area.

We believe that PA should increase with more time after bariatric surgery, since the change in the body composition of the operated patient will reflect an improvement in body mass distribution and, consequently, less inflammatory process. An important point is what form of protein supplementation should be used with these patients after surgery in order to minimize the loss of muscle mass and thereby increase PA. With the data presented in this study, we suggest that PA may be a marker of the state of

body composition linked to the functionality of SMM.

ARTICLE HIGHLIGHTS

Research background

Obesity is a complex chronic inflammatory disease characterized by excessive accumulation of body fat. In obese people, a common comorbidity is non-alcoholic fatty liver disease (NAFLD). NAFLD is considered the most common liver disease in Western countries, affecting 90% of morbidly obese patients eligible for bariatric surgery. To evaluate these patients, bioimpedance (BIA) can be a good method for nutritional and prognostic evaluation, using phase angle (PA).

Research motivation

There are not enough studies to evaluate morbid obesity, its body composition (described by the BIA), and associated comorbidities, such as NAFLD. We believe that the body change resulting from bariatric surgery will reflect an improvement in NAFLD and can be measured by PA, since it reflects cellular integrity and functionality by measuring, through an electrical current, the values of resistance and reactance of the membrane of these cells, with skeletal muscle being a conductor of electrical current and the opposite occurring with fat mass.

Research objectives

The aim of this study was to analyze the behavior of PA in the postoperative period of bariatric surgery, correlating it with changes in body composition and improvement of liver disease.

Research methods

This was a retrospective cohort study that analyzed the medical records of 727 patients undergoing bariatric surgery in a referral center of a teaching hospital in the south of Brazil. For convenience, the sample was carried out from July 2015 to July 2017. The data obtained were related to the protocol for routine pre- and postoperative care at the service's outpatient clinic. Quantitative and categorical variables analyses were performed to assess the association between PA, NAFLD, and body composition before and after bariatric surgery.

Research results

We analyzed 727 patients' medical records, and 379 patients were selected for having all preoperative data. Regarding PA, 169 patients were analyzed, and 33 patients had liver biopsy pre- and postoperatively with NAFLD information. The PA showed a significant reduction in the postoperative period as well as body composition data. Regarding liver disease, all patients presented a reduction in the degrees and stages of liver disease in the postoperative period, and some had no degree of liver disease.

Research conclusions

The PA decreased after bariatric surgery, with a direct correlation with weight loss and changes in body composition, and it did not correlate with the improvement of NAFLD. With the data presented in this study, we suggest that PA may be a marker of the state of body composition linked to the functionality of skeletal muscle mass.

Research perspectives

Performing large scale prospective studies with long-term follow-up are needed to verify if PA increases with more time after bariatric surgery, since the change in the body composition of the operated patient will reflect an improvement in body mass distribution and, consequently, less inflammatory process.

REFERENCES

- 1 Associação Brasileira para o Estudo da Obesidade e da Síndrome Metabólica (ABESO). Diretrizes brasileiras de obesidade, 2019. Available from: <https://abeso.org.br/wp-content/uploads/2019/12/Diretrizes-Download-Diretrizes-Brasileiras-de-Obesidade-2016.pdf>
- 2 de Moraes TS. Nutritional intervention in the treatment of patients obesity. Revista Brasileira de

- Obesidade, Nutrição e Emagrecimento 2007; 1: 38-47. Available from: <http://www.rbone.com.br/index.php/rbone/article/view/27/25>
- 3 **Prado WL**, Lofrano MC, Oyama LM, Dâmaso AR. Obesity and inflammatory adipokines: Practical implications for exercise prescription. *Rev Bras Med Esporte* 2009; **15**: 378-383 [DOI: [10.1590/S1517-86922009000600012](https://doi.org/10.1590/S1517-86922009000600012)]
 - 4 Obesity: preventing and managing the global epidemic. Report of a WHO consultation. *World Health Organ Tech Rep Ser* 2000; **894**: i-xii, 1 [PMID: [11234459](https://pubmed.ncbi.nlm.nih.gov/11234459/) DOI: [10.1002/jps.3080150106](https://doi.org/10.1002/jps.3080150106)]
 - 5 **Conway B**, Rene A. Obesity as a disease: no lightweight matter. *Obes Rev* 2004; **5**: 145-151 [PMID: [15245383](https://pubmed.ncbi.nlm.nih.gov/15245383/) DOI: [10.1111/j.1467-789X.2004.00144.x](https://doi.org/10.1111/j.1467-789X.2004.00144.x)]
 - 6 **Brasil**. Ministério da Saúde. VIGITEL Brasil 2018 [cited 2019 Nov 5]. Available from: <https://portal.arquivos2.saude.gov.br/images/pdf/2019/julho/25/vigitel-brasil-2018.pdf>
 - 7 **European Association for the Study of the Liver (EASL)**. European Association for the Study of Diabetes (EASD); European Association for the Study of Obesity (EASO). EASL-EASD-EASO Clinical Practice Guidelines for the management of non-alcoholic fatty liver disease. *J Hepatol* 2016; **64**: 1388-1402 [PMID: [27062661](https://pubmed.ncbi.nlm.nih.gov/27062661/) DOI: [10.1016/j.jhep.2015.11.004](https://doi.org/10.1016/j.jhep.2015.11.004)]
 - 8 **Marchesini G**, Petta S, Dalle Grave R. Diet, weight loss, and liver health in nonalcoholic fatty liver disease: Pathophysiology, evidence, and practice. *Hepatology* 2016; **63**: 2032-2043 [PMID: [26663351](https://pubmed.ncbi.nlm.nih.gov/26663351/) DOI: [10.1002/hep.28392](https://doi.org/10.1002/hep.28392)]
 - 9 **Losekann A**, Weston AC, Carli LA, Espindola MB, Pioner SR, Coral GP. Nonalcoholic fatty liver disease in severe obese patients, subjected to bariatric surgery. *Arq Gastroenterol* 2013; **50**: 285-289 [PMID: [24474231](https://pubmed.ncbi.nlm.nih.gov/24474231/) DOI: [10.1590/S0004-28032013000400009](https://doi.org/10.1590/S0004-28032013000400009)]
 - 10 **Chalasanani N**, Younossi Z, Lavine JE, Diehl AM, Brunt EM, Cusi K, Charlton M, Sanyal AJ. The diagnosis and management of non-alcoholic fatty liver disease: practice Guideline by the American Association for the Study of Liver Diseases, American College of Gastroenterology, and the American Gastroenterological Association. *Hepatology* 2012; **55**: 2005-2023 [PMID: [22488764](https://pubmed.ncbi.nlm.nih.gov/22488764/) DOI: [10.1002/hep.25762](https://doi.org/10.1002/hep.25762)]
 - 11 **Tovo CV**, Fernandes SA, Buss C, de Mattos AA. Sarcopenia and non-alcoholic fatty liver disease: Is there a relationship? *World J Hepatol* 2017; **9**: 326-332 [PMID: [28293382](https://pubmed.ncbi.nlm.nih.gov/28293382/) DOI: [10.4254/wjvh.v9.i6.326](https://doi.org/10.4254/wjvh.v9.i6.326)]
 - 12 **Losekann A**, Weston AC, de Mattos AA, Tovo CV, de Carli LA, Espindola MB, Pioner SR, Coral GP. Non-Alcoholic Steatohepatitis (NASH): Risk Factors in Morbidly Obese Patients. *Int J Mol Sci* 2015; **16**: 25552-25559 [PMID: [26512661](https://pubmed.ncbi.nlm.nih.gov/26512661/) DOI: [10.3390/ijms161025552](https://doi.org/10.3390/ijms161025552)]
 - 13 **Picot J**, Jones J, Colquitt JL, Gospodarevskaya E, Loveman E, Baxter L, Clegg AJ. The clinical effectiveness and cost-effectiveness of bariatric (weight loss) surgery for obesity: a systematic review and economic evaluation. *Health Technol Assess* 2009; **13**: 1-190, 215 [PMID: [19726018](https://pubmed.ncbi.nlm.nih.gov/19726018/) DOI: [10.3310/hta13410](https://doi.org/10.3310/hta13410)]
 - 14 **Cummings DE**, Overduin J, Foster-Schubert KE. Gastric bypass for obesity: mechanisms of weight loss and diabetes resolution. *J Clin Endocrinol Metab* 2004; **89**: 2608-2615 [PMID: [15181031](https://pubmed.ncbi.nlm.nih.gov/15181031/) DOI: [10.1210/jc.2004-0433](https://doi.org/10.1210/jc.2004-0433)]
 - 15 **Bordalo LA**, Teixeira TF, Bressan J, Mourão DM. [Bariatric surgery: how and why to supplement]. *Rev Assoc Med Bras (1992)* 2011; **57**: 113-120 [PMID: [21390468](https://pubmed.ncbi.nlm.nih.gov/21390468/) DOI: [10.1590/s0104-42302011000100025](https://doi.org/10.1590/s0104-42302011000100025)]
 - 16 **Rezende F**, Rosado L, Franceschini S, Rosado G, Ribeiro R, Marins JC. [Critical revision of the available methods for evaluate the body composition in population-based and clinical studies]. *Arch Latinoam Nutr* 2007; **57**: 327-334 [PMID: [18524316](https://pubmed.ncbi.nlm.nih.gov/18524316/)]
 - 17 **Buchholz AC**, Bartok C, Schoeller DA. The validity of bioelectrical impedance models in clinical populations. *Nutr Clin Pract* 2004; **19**: 433-446 [PMID: [16215137](https://pubmed.ncbi.nlm.nih.gov/16215137/) DOI: [10.1177/0115426504019005433](https://doi.org/10.1177/0115426504019005433)]
 - 18 **Coppini LZ**, Waitzberg DL, Campos AC. Limitations and validation of bioelectrical impedance analysis in morbidly obese patients. *Curr Opin Clin Nutr Metab Care* 2005; **8**: 329-332 [PMID: [15809537](https://pubmed.ncbi.nlm.nih.gov/15809537/) DOI: [10.1097/01.mco.0000165013.54696.64](https://doi.org/10.1097/01.mco.0000165013.54696.64)]
 - 19 **Marroni CA**, Miranda D, Boemeke L, Fernandes SA. Phase Angle Bioelectrical Impedance Analysis (BIA) as a Biomarker Tool for Liver Disease. In: Patel V, Preedy V. (eds) Biomarkers in Liver Disease. Biomarkers in Disease: Methods, Discoveries and Applications. Springer, Dordrecht. 2017: 735-751 [DOI: [10.1007/978-94-007-7675-3_43](https://doi.org/10.1007/978-94-007-7675-3_43)]
 - 20 **Fernandes SA**, de Mattos AA, Tovo CV, Marroni CA. Nutritional evaluation in cirrhosis: Emphasis on the phase angle. *World J Hepatol* 2016; **8**: 1205-1211 [PMID: [27803765](https://pubmed.ncbi.nlm.nih.gov/27803765/) DOI: [10.4254/wjvh.v8.i29.1205](https://doi.org/10.4254/wjvh.v8.i29.1205)]
 - 21 **Boemeke L**, Raimundo FV, Bopp M, Leonhardt LR, Fernandes SA, Marroni CA. The correlation of neck circumference and insulin resistance in NAFLD patients. *Arq Gastroenterol* 2019; **56**: 28-33 [PMID: [31141077](https://pubmed.ncbi.nlm.nih.gov/31141077/) DOI: [10.1590/S0004-2803.201900000-06](https://doi.org/10.1590/S0004-2803.201900000-06)]
 - 22 **Faria SL**, Faria OP, Cardeal MD, Ito MK. Validation study of multi-frequency bioelectrical impedance with dual-energy X-ray absorptiometry among obese patients. *Obes Surg* 2014; **24**: 1476-1480 [PMID: [24464546](https://pubmed.ncbi.nlm.nih.gov/24464546/) DOI: [10.1007/s11695-014-1190-5](https://doi.org/10.1007/s11695-014-1190-5)]
 - 23 **Baumgartner RN**, Heymsfield SB, Lichtman S, Wang J, Pierson RN Jr. Body composition in elderly people: effect of criterion estimates on predictive equations. *Am J Clin Nutr* 1991; **53**: 1345-1353 [PMID: [2035461](https://pubmed.ncbi.nlm.nih.gov/2035461/) DOI: [10.1093/ajcn/53.6.1345](https://doi.org/10.1093/ajcn/53.6.1345)]
 - 24 **Máttar JA**. Application of total body bioimpedance to the critically ill patient. Brazilian Group for Bioimpedance Study. *New Horiz* 1996; **4**: 493-503 [PMID: [8968982](https://pubmed.ncbi.nlm.nih.gov/8968982/)]
 - 25 **Kyle UG**, Bosaeus I, De Lorenzo AD, Deurenberg P, Elia M, Manuel Gómez J, Lilienthal Heitmann B, Kent-Smith L, Melchior JC, Pirlich M, Scharfetter H, M W J Schols A, Pichard C; ESPEN. Bioelectrical impedance analysis-part II: utilization in clinical practice. *Clin Nutr* 2004; **23**: 1430-1453 [PMID: [15556267](https://pubmed.ncbi.nlm.nih.gov/15556267/) DOI: [10.1016/j.clnu.2004.09.012](https://doi.org/10.1016/j.clnu.2004.09.012)]
 - 26 **Kleiner DE**, Brunt EM, Van Natta M, Behling C, Contos MJ, Cummings OW, Ferrell LD, Liu YC, Torbenson MS, Unalp-Arida A, Yeh M, McCullough AJ, Sanyal AJ; Nonalcoholic Steatohepatitis Clinical Research Network. Design and validation of a histological scoring system for nonalcoholic fatty liver disease. *Hepatology* 2005; **41**: 1313-1321 [PMID: [15915461](https://pubmed.ncbi.nlm.nih.gov/15915461/) DOI: [10.1002/hep.20701](https://doi.org/10.1002/hep.20701)]
 - 27 **Mottin CC**, Moretto M, Padoin AV, Kupski C, Swarowsky AM, Glock L, Duval V, da Silva JB. Histological behavior of hepatic steatosis in morbidly obese patients after weight loss induced by bariatric

- surgery. *Obes Surg* 2005; **15**: 788-793 [PMID: [15978148](#) DOI: [10.1381/0960892054222830](#)]
- 28 **Mattar SG**, Velcu LM, Rabinovitz M, Demetris AJ, Krasinskas AM, Barinas-Mitchell E, Eid GM, Ramanathan R, Taylor DS, Schauer PR. Surgically-induced weight loss significantly improves nonalcoholic fatty liver disease and the metabolic syndrome. *Ann Surg* 2005; **242**: 610-617; discussion 618 [PMID: [16192822](#) DOI: [10.1097/01.sla.0000179652.07502.3f](#)]
- 29 **Liu X**, Lazenby AJ, Clements RH, Jhala N, Abrams GA. Resolution of nonalcoholic steatohepatitis after gastric bypass surgery. *Obes Surg* 2007; **17**: 486-492 [PMID: [17608261](#) DOI: [10.1007/s11695-007-9086-2](#)]
- 30 **Clark JM**, Alkhuraishi AR, Solga SF, Alli P, Diehl AM, Magnuson TH. Roux-en-Y gastric bypass improves liver histology in patients with non-alcoholic fatty liver disease. *Obes Res* 2005; **13**: 1180-1186 [PMID: [16076987](#) DOI: [10.1038/oby.2005.140](#)]
- 31 **Moretto M**, Kupski C, da Silva VD, Padoin AV, Mottin CC. Effect of bariatric surgery on liver fibrosis. *Obes Surg* 2012; **22**: 1044-1049 [PMID: [22108808](#) DOI: [10.1007/s11695-011-0559-y](#)]
- 32 **Stefan N**, Häring HU, Cusi K. Non-alcoholic fatty liver disease: causes, diagnosis, cardiometabolic consequences, and treatment strategies. *Lancet Diabetes Endocrinol* 2019; **7**: 313-324 [PMID: [30174213](#) DOI: [10.1016/S2213-8587\(18\)30154-2](#)]
- 33 **Llames L**, Baldomero V, Iglesias ML, Rodota LP. [Values of the phase angle by bioelectrical impedance; nutritional status and prognostic value]. *Nutr Hosp* 2013; **28**: 286-295 [PMID: [23822677](#) DOI: [10.3305/nh.2013.28.2.6306](#)]
- 34 **Perrone F**, Bianciardi E, Benavoli D, Tognoni V, Niolu C, Siracusano A, Gaspari AL, Gentileschi P. Gender Influence on Long-Term Weight Loss and Comorbidities After Laparoscopic Sleeve Gastrectomy and Roux-en-Y Gastric Bypass: a Prospective Study With a 5-Year Follow-up. *Obes Surg* 2016; **26**: 276-281 [PMID: [26033435](#) DOI: [10.1007/s11695-015-1746-z](#)]
- 35 **Hartwig TW**, dos Santos FAI, González MC, Rombaldi AJ. Bariatric surgery and body composition of adults. *Rev Bras Cineantropom Desempenho Hum* 2013; **15**: 686-694 [DOI: [10.5007/1980-0037.2013v15n6p686](#)]
- 36 **de Paris FGC**, Padoin AV, Mottin CC, de Paris MF. Assessment of Changes in Body Composition During the First Postoperative Year After Bariatric Surgery. *Obes Surg* 2019; **29**: 3054-3061 [PMID: [31254214](#) DOI: [10.1007/s11695-019-03980-8](#)]
- 37 **Perumpail BJ**, Khan MA, Yoo ER, Cholankeril G, Kim D, Ahmed A. Clinical epidemiology and disease burden of nonalcoholic fatty liver disease. *World J Gastroenterol* 2017; **23**: 8263-8276 [PMID: [29307986](#) DOI: [10.3748/wjg.v23.i47.8263](#)]
- 38 **Younossi ZM**, Koenig AB, Abdelatif D, Fazel Y, Henry L, Wymer M. Global epidemiology of nonalcoholic fatty liver disease-Meta-analytic assessment of prevalence, incidence, and outcomes. *Hepatology* 2016; **64**: 73-84 [PMID: [26707365](#) DOI: [10.1002/hep.28431](#)]
- 39 **Gonzalez MC**, Heymsfield SB. Bioelectrical impedance analysis for diagnosing sarcopenia and cachexia: what are we really estimating? *J Cachexia Sarcopenia Muscle* 2017; **8**: 187-189 [PMID: [28145079](#) DOI: [10.1002/jcsm.12159](#)]
- 40 **Basile C**, Della-Morte D, Cacciatore F, Gargiulo G, Galizia G, Roselli M, Curcio F, Bonaduce D, Abete P. Phase angle as bioelectrical marker to identify elderly patients at risk of sarcopenia. *Exp Gerontol* 2014; **58**: 43-46 [PMID: [25034911](#) DOI: [10.1016/j.exger.2014.07.009](#)]
- 41 **Kilic MK**, Kizilarslanoglu MC, Arik G, Bolayir B, Kara O, Dogan Varan H, Sumer F, Kuyumcu ME, Halil M, Ulger Z. Association of Bioelectrical Impedance Analysis-Derived Phase Angle and Sarcopenia in Older Adults. *Nutr Clin Pract* 2017; **32**: 103-109 [PMID: [27590205](#) DOI: [10.1177/0884533616664503](#)]
- 42 **Stobäus N**, Pirlich M, Valentini L, Schulzke JD, Norman K. Determinants of bioelectrical phase angle in disease. *Br J Nutr* 2012; **107**: 1217-1220 [PMID: [22309898](#) DOI: [10.1017/S0007114511004028](#)]
- 43 **Bosy-Westphal A**, Danielzik S, Dörhöfer RP, Later W, Wiese S, Müller MJ. Phase angle from bioelectrical impedance analysis: population reference values by age, sex, and body mass index. *JPEN J Parenter Enteral Nutr* 2006; **30**: 309-316 [PMID: [16804128](#) DOI: [10.1177/0148607106030004309](#)]
- 44 **Barbosa-Silva MC**, Barros AJ. Bioelectric impedance and individual characteristics as prognostic factors for post-operative complications. *Clin Nutr* 2005; **24**: 830-838 [PMID: [15975694](#) DOI: [10.1016/j.clnu.2005.05.005](#)]
- 45 **Baumgartner RN**, Chumlea WC, Roche AF. Bioelectric impedance phase angle and body composition. *Am J Clin Nutr* 1988; **48**: 16-23 [PMID: [3389323](#) DOI: [10.1093/ajcn/48.1.16](#)]
- 46 **Peres WA**, Lento DF, Baluz K, Ramalho A. Phase angle as a nutritional evaluation tool in all stages of chronic liver disease. *Nutr Hosp* 2012; **27**: 2072-2078 [PMID: [23588459](#) DOI: [10.3305/nh.2012.27.6.6015](#)]
- 47 **Norman K**, Stobäus N, Pirlich M, Bosy-Westphal A. Bioelectrical phase angle and impedance vector analysis--clinical relevance and applicability of impedance parameters. *Clin Nutr* 2012; **31**: 854-861 [PMID: [22698802](#) DOI: [10.1016/j.clnu.2012.05.008](#)]
- 48 **Vassilev G**, Hasenberg T, Krammer J, Kienle P, Ronellenfisch U, Otto M. The Phase Angle of the Bioelectrical Impedance Analysis as Predictor of Post-Bariatric Weight Loss Outcome. *Obes Surg* 2017; **27**: 665-669 [PMID: [27465938](#) DOI: [10.1007/s11695-016-2315-9](#)]
- 49 **Koehler KB**, Moraes RAG, Rodrigues JB, Portela BSM, Miguel GPS, Pedrosa RG, Haraguchi FK. Bioimpedance phase angle is associated with serum transthyretin but not with prognostic inflammatory and nutritional index during follow-up of women submitted to bariatric surgery. *Clin Nutr ESPEN* 2019; **33**: 183-187 [PMID: [31451259](#) DOI: [10.1016/j.clnesp.2019.05.018](#)]
- 50 **Cazzo E**, Pareja JC, Chaim EA. Nonalcoholic fatty liver disease and bariatric surgery: a comprehensive review. *Sao Paulo Med J* 2017; **135**: 277-295 [PMID: [28562737](#) DOI: [10.1590/1516-3180.2016.0306311216](#)]
- 51 **Barker KB**, Palekar NA, Bowers SP, Goldberg JE, Pulcini JP, Harrison SA. Non-alcoholic steatohepatitis: effect of Roux-en-Y gastric bypass surgery. *Am J Gastroenterol* 2006; **101**: 368-373 [PMID: [16454845](#) DOI: [10.1111/j.1572-0241.2006.00419.x](#)]
- 52 **Csendes A**, Smok G, Burgos AM. Histological findings in the liver before and after gastric bypass. *Obes Surg* 2006; **16**: 607-611 [PMID: [16687030](#) DOI: [10.1381/096089206776944904](#)]

- 53 **de Almeida SR**, Rocha PR, Sanches MD, Leite VH, da Silva RA, Diniz MT, Diniz Mde F, Rocha AL. Roux-en-Y gastric bypass improves the nonalcoholic steatohepatitis (NASH) of morbid obesity. *Obes Surg* 2006; **16**: 270-278 [PMID: [16545157](#) DOI: [10.1381/096089206776116462](#)]
- 54 **Silverman EM**, Sapala JA, Appelman HD. Regression of hepatic steatosis in morbidly obese persons after gastric bypass. *Am J Clin Pathol* 1995; **104**: 23-31 [PMID: [7611176](#) DOI: [10.1093/ajcp/104.1.23](#)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA
Telephone: +1-925-3991568
E-mail: bpgoffice@wjgnet.com
Help Desk: <https://www.f6publishing.com/helpdesk>
<https://www.wjgnet.com>

